

1 **Effect of crude glycerin supplementation via drinking water on feed intake, water intake**
2 **and productive performance of dairy ewes**

3
4
5 Felipe Santiago Santos, Hemilly Cristina Menezes de Sá, Luciano Soares de Lima, Tássia
6 Ludmila Teles Martins, Flávio Augusto Pereira Alvarenga, José André Junior, Luciana Freitas
7 Guedes, Leonardo de Rago Nery Alves, Joana Palhares Campolina, Gilberto de Lima Macedo
8 Júnior and Iran Borges

9
10 SUPPLEMENTARY FILE

11
12 Supplementary Table S1. Ingredients and chemical composition of experimental diets

| | |
|--------------------------------------|-------|
| Ingredient (g/kg DM) | |
| Tifton hay | 503.3 |
| Ground corn | 315.1 |
| Soybean meal | 162.2 |
| Mineral Supplement ¹ | 10.0 |
| Salt | 5.0 |
| Limestone | 3.46 |
| Dicalcium phosphate | 0.95 |
| Chemical composition | |
| Dry matter (g/kg) | 895.6 |
| Crude Protein (g/kg DM) | 155.7 |
| Neutral detergent fiber (g/kg DM) | 420.0 |
| Acid detergent fiber (g/kg DM) | 210.0 |
| Calcium (g/kg DM) | 4.3 |
| Phosphorus (g/kg DM) | 3.7 |
| Total digestible nutrients (g/kg DM) | 695.3 |

13 ¹Contained (per kg, as-fed basis): Ca 110 g, P 87 g, Na 147 g, Zn 3,800 mg, Mn 2,000 mg, Co
14 15 mg, Co 590 mg, S 18 g, Se 20 mg, I 50 mg, Cr 20 mg, Mo 300 mg.

15