

Supplementary Materials for
IS THE RISK OF BANK VOLE INFECTION WITH CESTODES RELATED
TO THE POPULATION DYNAMICS REGIME?

Yulia A. Davydova*, Ivan A. Kshnyasev

Institute of Plant and Animal Ecology, Ural Branch of the Russian Academy of Sciences,
8 Marta Str. 202, 620144, Yekaterinburg, Russia

Corresponding author: Yu. A. Davydova; Emails: davydova@ipae.uran.ru

Table S1. The number of non-infected (0) and infected (1) *Clethrionomys glareolus* with cestode larvae in different population dynamics regimes

Regime of dynamics	Phase of population cycle	Year	Parasite infection		Sum
			0	1	
Cyclic (3-year cycle)	Peak	1995	126	3	129
	Trough	1996	14	0	14
	Increase	1997	159	0	159
	Peak	1998	74	1	75
	Trough	1999	32	0	32
	Increase	2000	66	0	66
	Peak	2001	208	11	219
	Trough	2002	11	0	11
	Increase	2003	136	4	140
	Peak	2004	116	13	129
	Trough	2005	10	0	10
Increase	2006	69	19	88	
Noncyclic (quasi-2-year cycle)	–	2007	61	0	61
	–	2008	70	9	79
	–	2009	36	1	37
	–	2010	68	2	70
	–	2011	25	2	27
	–	2012	84	5	89
	–	2013	29	5	34
	–	2014	83	1	84
	–	2015	60	4	64
–	2016	34	3	37	
Cyclic (3-year cycle)	Peak	2017	88	2	90
	Trough	2018	8	1	9
	Increase	2019	35	1	36
	Peak	2020	136	8	144
	Trough	2021	3	2	5
Totals			1841	97	1938