

Online supplement, estimates of psychiatric bed needs


Supplement Table 1:

Supplement Table 1. Number of studies per country, categorized by World Bank Income group of the country.


Income group	Country	Studies
High-income	United Kingdom	24
	USA	12
	Australia	8
	Canada	6
	Germany	2
	Denmark, Ireland, Japan, Netherlands, New Zealand, Norway, Spain, Switzerland*	1
Low- and middle-income	Brazil	2
	South Africa	1
	International	3
Total		66

*One study for each country.


Supplement Figure 1: Need estimates of psychiatric beds with unspecified length of stay per 100 000 population




Supplement Figure 2: Need estimates of psychiatric short-stay beds per 100 000 population




Supplement Figure 3: Need estimates of psychiatric long-stay beds per 100 000 population




Supplement Figure 4: Trendlines of need estimates for psychiatric beds per 100 000 population over time.




Supplementary Figure 5: Trendlines of need estimates for the psychiatric bed prevalence from 2000 onwards.



Supplement Figure 6: Prevalence ratios between need estimates and actually available psychiatric beds by length of stay.



Supplementary Figure 7: Linear trendlines for prevalence ratio between estimates of psychiatric bed needs and actual availability over time (from 1990 onwards) on a semi logarithmic scale. Prevalence ratios smaller than 1 indicate recommendations to remove, and greater than 1, recommendations to increase psychiatric bed numbers. Lines are curved due to the logarithmic scale.



Supplementary Table 2: Quality appraisal of the scientific literature reporting need estimates for psychiatric beds.

	Supplement Table 2. Quality appraisal of publications.							
	<i>Author (year of publication)</i>	<i>Published in indexed journal with peer review</i>	<i>Recommendation based on original epidemiological data</i>	<i>Primary aim of the study was the recommendation of bed numbers</i>	<i>Clear definition of length of stay</i>	<i>Provided as prevalences of beds per population</i>	<i>Provided as a single number (vs range)</i>	<i>Definition of recommendation as "minimal"^a</i>
1	Strömgren E (1958)	X	X	X		X	X	X
2	Tooth GC & Brooke E (1961)	X	X	X	X	X	X	
3	Gore et al (1964)	X	X	X	X			
4	Richman & Kennedy (1965)	X	X	X	X	X	X	
5	Ohio Department of Mental Hygiene and Correction (1965)			X	X	X	X	
6	Maryland Board of Health and Mental Hygiene (1965)			X		X	X	
7	Valdés AJ (1966)	X		X		X	X	
8	Orwin A (1967)	X	X	X	X	X	X	X
9	California Department of Mental Hygiene (1968)			X	X	X		
10	Baker et al (1969)	X	X	X	X	X	X	X
11	Oldham AJ (1969)	X	X	X			X	
12	Hoening J & Hamilton MW (1969)		X	X	X	X	X	X
13	Crane & Gorwic (1970)		X	X	X	X	X	
14	Cross et al (1970)	X	X	X	X	X	X	
15	Hailey (1971)	X	X	X	X	X	X	
16	Department Health and Social Security (1971)			X	X	X	X	X
17	Mezey & Syed (1972)	X	X	X	X	X	X	
18	Lipscomb CF (1972)	X	X	X	X	X		
19	Lawrence M (1972)				X	X	X	
20	Lawson JS (1972)	X		X		X	X	
21	Brown BM (1973)		X	X		X		
22	Muller C (1973)	X		X		X	X	
23	Fryers T. (1974 ^b)	X	X	X	X	X	X	
24	UK Department Health and Social Security (1975)			X	X	X	X	
25	Straathof LJ (1976)	X	X	X	X	X	X	
26	Robertson G (1981)	X	X	X	X	X	X	
27	Hirsch SR (1983)	X	X	X				X
28	McGreadie et al (1983)	X	X	X	X	X	X	
29	Goplerud EN (1984)	X	X	X	X	X		
30	Richman et al (1984)	X	X	X	X	X	X	
31	Gudeman & Shore (1984)	X	X	X	X	X	X	
32	Trant et al (1984)		X	X	X	X	X	X
33	Häfner H (1987)	X	X	X	X	X		X
34	Hirsch SR (1987)	X	X	X	X	X		X
35	Andrews et al (1990)	X	X			X	X	X
36	Holloway et al (1992)	X	X		X	X	X	
37	Strathdee & Thornicroft (1992)		X	X	X	X		
38	Wing JK (1992)		X	X	X	X		
39	Pridmore & Jones (1992)	X	X	X	X	X	X	X
40	Rosenman S (1995)	X	X	X	X	X	X	
41	World Health Organisation (1996)			X		X	X	
42	Trieman & Leff (1996)	X	X	X	X	X		
43	Gordon A (1997)			X		X		X
44	Ramsay et al (1997)		X	X	X	X		
45	Dillard et al (1998)		X	X	X	X	X	
46	Disley et al (1998)		X	X	X	X	X	X
47	Audini et al (1999)	X	X	X	X	X		X
48	Gebhardt (2002)	X	X	X	X		X	X
49	Lasage et al (2002)	X	X	X	X	X	X	X
50	Lund & Flisher (2006)	X	X	X	X	X	X	
51	Boardman & Parsonage (2007)		X	X	X	X	X	
52	Leite Gastal et al (2007)	X				X	X	
53	Fuller Torrey et al (2008)			X		X		X
54	Miklebust et al (2008)	X	X	X		X	X	X
55	Candiago et al (2010)	X	X					X
56	Tyler P (2011)	½ ^b		X			X	X
57	Harris et al (2012)	X	X	X	X	X	X	
58	La et al (2016)	X	X	X				X
59	Allison et al (2018)	X		X		X	X	X
60	Benjamin et al (2018)	½ ^b		X	X		X	X
61	Victoria's Mental Health System (2019)		X	X	X	X	X	
62	Hudson (2020)	X	X	X		X	X	X
63	Hudson (2021)	X	X	X		X	X	X
64	McBain et al (2021)		X	X	X	X	X	X
65	Mundt et al (2022)	X		X	X	X	X	X

^a "Minimal" considers all reports that refer to the recommendation as "minimal", "sufficient", "at least", "necessary", "bedrock" or "required".

^b Indexed journal, but format that does not typically undergo peer review, like correspondence