

Supplemental Appendix - Overview of the studies

Reference	Design	Sample size	Type of PS	Food format	Dose of PS ¹	Relative change in LDL-cholesterol	
						Effect (%)	SE
AbuMweis et al. 2006 #1 ²	x	30	sterol	margarine	1.7	-1.4	3.1
AbuMweis et al. 2006 #2	x	30	sterol	margarine	1.7	-1.4	3.2
Algorta-Pineda et al. 2005	p	32	stanol	yoghurt drink	2.0	-8.4	3.9
Alhassan et al. 2006	p	26	stanol	low-fat margarine	3.0	-20.4	6.6
Andersson et al. 1999	p	40	stanol	low-fat margarine	1.9	-7.2	2.9
Athyros et al. 2011	p	100	stanol	margarine	2.0	-13.8	1.3
Banuls et al. 2010	p	40	sterol	low-fat milk	2.0	-8.1	2.8
Banuls et al. 2011	p	75	sterol	low-fat milk	2.0	-9.9	2.8
Beer et al 2000 #1	p	47	mix	low-fat milk	0.9	-7.4	3.5
Beer et al 2000 #2	p	52	mix	low-fat milk	1.8	-8.6	3.3
Beer et al 2000 #3	p	47	mix	low-fat milk	3.6	-13.2	3.4
Blair et al. 2000	p	141	stanol	margarine	2.9	-9.0	1.8
Blomqvist et al. 1993	p	66	stanol	mayonnaise	3.4	-9.9	3.3
Cater et al. 2005 study 1 ³ #1	x	8	stanol	margarine	2.0	-12.3	3.3
Cater et al. 2005 study 1 #2	x	8	stanol	margarine	3.0	-13.0	2.6
Cater et al. 2005 study 1 #3	x	8	stanol	margarine	4.0	-13.6	2.3
Cater et al. 2005 study 2	x	13	stanol	margarine	3.0	-13.0	2.3
Cater et al. 2005 study 3	x	10	stanol	margarine	3.0	-14.9	3.3
Chen et al. 2009	x	22	sterol	dressing and margarine	3.3	-12.4	1.9
Christiansen et al. 2001 #1	p	92	sterol	margarine	1.5	-6.4	1.9
Christiansen et al. 2001 #2	p	88	sterol	margarine	3.0	-9.1	2.4
Cleghorn et al. 2003	x	50	sterol	margarine	2.1	-7.2	1.7
Clifton et al. 2004 #1	x	36	sterol	bread	1.6	-9.8	1.4
Clifton et al. 2004 #2	x	40	sterol	milk	1.6	-12.4	1.3
Clifton et al. 2004 #3	x	58	sterol	cereals	1.6	-5.6	1.3
Clifton et al. 2004 #4	x	40	sterol	yoghurt	1.6	-9.8	1.2
Clifton et al. 2008 #1	p	76	sterol	low-fat spread	1.6	-9.1	3.4
Clifton et al. 2008 #2	p	78	sterol	low-fat spread	1.6	-11.4	3.3
Clifton et al. 2008 #3	p	75	sterol	low-fat spread	1.6	-7.3	3.4
Colgan et al. 2004	x	48	sterol	low-fat margarine	1.3	-2.9	2.4
Davidson et al. 2001 #1	p	38	sterol	low-fat spread	3.0	-4.3	5.0
Davidson et al. 2001 #2	p	37	sterol	dressing	6.0	-3.9	4.5
Davidson et al. 2001 #3	p	40	sterol	low-fat spread and dressing	9.0	-12.2	4.9
de Graaf et al. 2002	p	62	mix	chocolate	1.8	-11.1	3.1
de Jong et al. 2008a #1	p	26	sterol	low-fat margarine	2.5	-8.2	5.6
de Jong et al. 2008a #2	p	26	stanol	low-fat margarine	2.5	-12.8	5.1
de Jong et al. 2008b #1	p	35	sterol	low-fat margarine	2.5	-12.4	4.3
de Jong et al. 2008b #2	p	36	stanol	low-fat margarine	2.5	-9.5	4.3
Devaraj et al. 2004	p	72	sterol	orange juice	2.0	-11.8	2.5
Devaraj et al. 2006	p	72	sterol	orange juice	2.0	2.1	3.7
Doornbos et al. 2006 #1	p	71	sterol	low-fat yoghurt drink	3.2	-9.5	2.2

Reference	Design	Sample size	Type of PS	Food format	Dose of PS ¹	Relative change in LDL-cholesterol	
						Effect (%)	SE
Doornbos et al. 2006 #2	p	71	sterol	low-fat yoghurt drink	2.8	-9.3	2.2
Eady et al. 2011	x	39	sterol	spread	1.6	-5.6	1.7
Earnest et al. 2007	p	54	sterol	capsules	1.6	-9.9	3.4
Escuriol et al. 2010	x	44	sterol	milk	2.0	-4.1	2.0
Fuentes et al. 2008 #1	x	30	sterol	low-fat margarine	2.0	-7.7	2.7
Fuentes et al. 2008 #2	x	30	sterol	low-fat margarine	2.0	-3.3	3.0
Geelen et al. 2002 study 1	x	31	sterol	low-fat margarine	3.2	-9.9	2.6
Geelen et al. 2002 study 2	x	57	sterol	low-fat margarine	3.2	-12.4	2.2
Goldberg et al. 2006	p	26	stanol	tablets	1.8	-9.1	4.5
Goncalves et al. 2007	p	34	sterol	milk	2.0	4.0	4.8
Gylling & Miettinen 1994	x	11	stanol	margarine	3.0	-9.3	2.8
Gylling & Miettinen 1996 #1	x	8	stanol	margarine	3.0	-14.4	2.1
Gylling & Miettinen 1996 #2	x	8	stanol	margarine	3.0	-9.7	3.9
Gylling & Miettinen 1999	x	21	stanol	butter	2.4	-12.0	2.0
Gylling et al. 1997	x	22	stanol	margarine	3.0	-14.5	2.9
Gylling et al. 2010	p	49	stanol	margarine and oat-based drink	8.9	-17.4	2.6
Hallikainen & Uusitupa 1999 #1	p	35	stanol	low-fat margarine	2.3	-14.0	3.0
Hallikainen & Uusitupa 1999 #2	p	37	stanol	low-fat margarine	2.2	-7.8	3.2
Hallikainen et al. 2000 #1	x	34	stanol	margarine	2.0	-12.7	2.2
Hallikainen et al. 2000 #2	x	34	sterol	margarine	2.0	-10.4	1.9
Hallikainen et al. 2008	p	19	stanol	margarine	2.2	-18.6	8.8
Hallikainen et al. 2011	p	24	stanol	margarine	3.2	-13.7	5.2
Hansel et al. 2007	p	194	sterol	low-fat fermented milk	1.6	-9.2	1.0
Hayes et al. 2004	x	7	sterol	tortilla chips	1.5	-15.3	5.9
Heggen et al. 2010 #1	x	59	sterol	low-fat margarine	2.0	-9.0	1.8
Heggen et al. 2010 #2	x	59	sterol	low-fat margarine	2.0	-8.2	1.6
Hendriks et al. 1999 #1	x	80	sterol	margarine	0.8	-6.2	1.8
Hendriks et al. 1999 #2	x	80	sterol	margarine	1.6	-9.2	1.8
Hendriks et al. 1999 #3	x	80	sterol	margarine	3.2	-9.8	1.8
Hendriks et al. 2003	p	185	sterol	low-fat spread	1.6	-4.3	2.4
Hernandez-Mijares et al. 2010	p	55	sterol	low-fat milk	2.0	-10.2	2.3
Hernandez-Mijares et al. 2011 study 1	p	24	sterol	low-fat milk	2.0	-0.5	3.8
Hernandez-Mijares et al. 2011 study 2	p	24	sterol	low-fat milk	2.0	-10.5	3.7
Hironaka et al. 2006 #1	p	101	sterol	vegetable/fruit juice	0.8	-6.7	2.1
Hironaka et al. 2006 #2	p	105	sterol	vegetable/fruit juice	1.6	-8.8	2.0
Homma et al. 2003 #1	p	67	stanol	low-fat spread	2.0	-8.9	1.9
Homma et al. 2003 #2	p	68	stanol	low-fat spread	3.0	-6.6	2.3
Houweling et al. 2009	x	82	sterol	low-fat margarine	2.0	-7.8	2.0
Hyun et al. 2005	p	51	stanol	low-fat yoghurt	2.0	-7.8	3.0
Ishizaki et al. 2003	p	55	sterol	mayonnaise	0.9	-8.2	2.4
Jakulj et al. 2005 #1	x	39	sterol	low-fat spread	2.0	-4.7	1.7
Jakulj et al. 2005 #2	x	39	sterol	low-fat spread	2.0	-3.5	2.3

Reference	Design	Sample size	Type of PS	Food format	Dose of PS ¹	Relative change in LDL-cholesterol	
						Effect (%)	SE
Jauhainen et al. 2006	p	67	stanol	low-fat cheese	2.0	-10.3	2.3
Jones et al. 1999	p	32	mix	margarine	1.7	-15.5	5.4
Jones et al. 2000 #1	x	15	sterol	low-fat margarine	1.8	-13.3	2.3
Jones et al. 2000 #2	x	15	stanol	low-fat margarine	1.8	-6.4	2.7
Jones et al. 2003	x	15	mix	non-fat beverage	1.8	-2.1	4.4
Judd et al. 2002	x	53	sterol	salad dressing	2.2	-10.1	0.7
Kassis et al. 2008	x	22	sterol	margarine	1.7	-4.5	1.3
Khandelwal et al. 2009 study 1	p	93	sterol	yoghurt drink	2.0	-3.3	2.9
Khandelwal et al. 2009 study 2	p	85	sterol	yoghurt drink	2.0	-5.0	2.8
Korpela et al. 2006 study 1	p	50	sterol	low-fat yoghurt	1.7	-7.7	3.6
Korpela et al. 2006 study 2	p	62	sterol	low-fat hard cheese	2.0	-11.2	3.4
Korpela et al. 2006 study 3	p	52	sterol	low-fat fresh cheese	2.0	-13.8	3.5
Kratz et al. 2007 #1	x	17	sterol	low-fat margarine	2.0	-4.0	4.2
Kratz et al. 2007#2	x	17	stanol	low-fat margarine	2.0	-9.1	2.7
Kurokawa et al. 2008a #1	p	35	sterol	dressing	0.5	-2.2	5.2
Kurokawa et al. 2008a #2	p	35	sterol	dressing	0.9	-6.7	5.2
Kurokawa et al. 2008a #3	p	34	sterol	dressing	1.3	-7.3	4.8
Kurokawa et al. 2008b	p	59	sterol	dressing	0.8	-5.2	1.7
Lagstrom et al. 2006	p	42	stanol	capsules	2.0	-7.0	3.2
Lau et al. 2005 study 1	x	14	sterol	margarine	1.8	-7.1	7.8
Lau et al. 2005 study 2	x	15	sterol	margarine	1.8	-8.4	3.8
Lee et al. 2003	p	81	sterol	low-fat spread	1.6	-8.1	2.4
Li et al. 2007 #1	p	201	sterol	milk tea powder	1.5	-2.5	1.6
Li et al. 2007 #2	p	199	sterol	milk tea powder	2.3	-3.4	1.6
Lin et al. 2011	x	21	sterol	beverage	2.5	-6.5	3.2
Lottenberg et al. 2003	x	60	sterol	margarine	1.7	-6.4	1.1
Madsen et al. 2007	x	46	sterol	low-fat margarine and low-fat milk	2.3	-7.7	2.2
Maki et al. 2001 #1	p	158	sterol	low-fat spread	1.1	-7.6	1.7
Maki et al. 2001 #2	p	118	sterol	low-fat spread	2.2	-8.1	1.9
Mannarino et al. 2009	p	116	sterol	low-fat fermented milk	1.6	-8.3	1.6
Matsuoka et al. 2004a	p	46	sterol	mayonnaise	0.8	0.2	3.1
Matsuoka et al. 2004b study 1 #1	p	16	sterol	mayonnaise	0.2	-5.3	4.9
Matsuoka et al. 2004b study 1 #2	p	19	sterol	mayonnaise	0.4	3.1	5.0
Matsuoka et al. 2004b study 1 #3	p	16	sterol	mayonnaise	0.6	-0.8	4.9
Matsuoka et al. 2004b study 1 #4	p	16	sterol	mayonnaise	0.8	-7.1	3.9
Matsuoka et al. 2004b study 2 #1	p	15	sterol	mayonnaise	0.8	-12.3	6.3
Matsuoka et al. 2004b study 2 #2	p	17	sterol	mayonnaise	1.6	-8.6	4.0
Matsuoka et al. 2004b study 2 #3	p	15	sterol	mayonnaise	2.4	-11.0	4.8
Matvienko et al. 2002	p	34	sterol	beef	2.7	-13.4	3.6
McPherson et al. 2005 study 1	p	25	stanol	tablets	1.3	-10.4	4.0
McPherson et al. 2005 study 2	p	27	stanol	capsules	1.0	-2.5	3.9
Mensink et al. 2002	p	60	stanol	yoghurt	3.0	-10.3	4.7

Reference	Design	Sample size	Type of PS	Food format	Dose of PS ¹	Relative change in LDL-cholesterol	
						Effect (%)	SE
Mensink et al. 2010 #1	p	46	stanol	margarine	2.8	-7.4	2.9
Mensink et al. 2010 #2	p	44	stanol	soy-based yoghurt	5.8	-11.9	3.2
Mensink et al. 2010 #3	p	47	stanol	margarine and soy-based yoghurt	8.7	-17.1	3.1
Miettinen and Vanhanen 1994 #1	p	17	sterol	mayonnaise	1.0	-6.2	3.8
Miettinen and Vanhanen 1994 #2	p	15	mix	mayonnaise	1.0	-2.6	4.1
Miettinen and Vanhanen 1994 #3	p	15	mix	mayonnaise	1.2	-7.7	3.7
Miettinen et al. 1995 #1	p	102	stanol	margarine	2.6	-11.1	2.1
Miettinen et al. 1995 #2	p	102	stanol	margarine	2.6	-12.5	2.2
Mussner et al. 2002	x	62	sterol	margarine	1.8	-6.5	1.4
Naumann et al. 2003 #1	x	42	mix	low-fat margarine	2.0	-6.0	3.1
Naumann et al. 2003 #2	x	42	mix	low-fat margarine	2.0	-6.7	3.0
Neil et al. 2001	p	62	sterol	spread	2.5	-14.2	3.3
Nestel et al. 2001	x	15	sterol	dairy spread	2.4	-7.9	2.5
Nigon et al. 2001	x	53	sterol	low-fat spread	1.6	-5.3	1.6
Niittynen et al. 2007 study 1	x	15	sterol	low-fat yoghurt drink	1.0	-4.2	3.3
Niittynen et al. 2007 study 2	p	26	sterol	low-fat yoghurt drink	2.0	-6.0	4.3
Noakes et al. 2002 study 1 #1	x	46	sterol	low-fat spread	2.3	-7.7	1.2
Noakes et al. 2002 study 1 #2	x	46	stanol	low-fat spread	2.5	-9.5	1.2
Noakes et al. 2002 study 2	x	35	sterol	spread	2.0	-9.6	1.5
Noakes et al. 2005 study 1 #1	x	39	sterol	margarine	2.0	-10.1	1.6
Noakes et al. 2005 study 1 #2	x	39	sterol	low-fat milk	2.0	-7.9	1.6
Noakes et al. 2005 study 1 #3	x	39	sterol	low-fat milk and margarine	4.0	-11.4	1.5
Noakes et al. 2005 study 2 #1	x	40	sterol	low-fat yoghurt	1.8	-6.1	1.6
Noakes et al. 2005 study 2 #2	x	40	stanol	low-fat yoghurt	1.7	-5.2	1.7
Ntanios et al. 2002	x	53	sterol	margarine	1.8	-9.1	1.6
Ooi et al. 2007	x	9	sterol	cereal and margarine	2.0	-6.4	7.8
Pelletier et al. 1995	x	12	sterol	butter	0.7	-15.2	3.1
Plana et al. 2008	p	83	sterol	low-fat fermented milk	1.6	-12.2	3.1
Plat and Mensink 2000 #1	p	78	stanol	margarine and shortening	3.8	-12.6	3.3
Plat and Mensink 2000 #2	p	76	stanol	margarine and shortening	4.0	-11.6	3.7
Plat et al. 2000 #1	x	39	stanol	margarine	2.5	-9.4	1.5
Plat et al. 2000 #2	x	39	stanol	margarine and cake/cookie	2.5	-10.4	1.9
Quilez et al. 2003	p	57	sterol	muffin and croissant	3.2	-14.7	4.5
Racette et al. 2010 #1	x	18	sterol	beverage	0.4	-5.0	2.1
Racette et al. 2010 #2	x	18	sterol	beverage	2.0	-8.9	2.3
Raitakari et al. 2008	p	190	stanol	low-fat margarine	2.0	-9.3	3.1
Rudkowska et al. 2008 #1	x	26	sterol	low-fat yoghurt	1.6	-2.3	1.8
Rudkowska et al. 2008 #2	x	26	sterol	low-fat yoghurt	1.6	-5.1	1.8
Ruiu et al. 2009	x	15	sterol	yoghurt drink	1.0	-4.6	2.2
Saito et al. 2006 #1	p	33	sterol	DAG-containing mayonnaise	0.3	-7.1	2.7
Saito et al. 2006 #2	p	33	sterol	DAG-containing mayonnaise	0.4	-5.9	3.2
Saito et al. 2006 #3	p	34	sterol	DAG-containing mayonnaise	0.5	-9.3	2.7

Reference	Design	Sample size	Type of PS	Food format	Dose of PS ¹	Relative change in LDL-cholesterol	
						Effect (%)	SE
Seki et al. 2003a	p	60	sterol	vegetable oil in bread	0.5	-2.3	2.4
Seki et al. 2003b	p	22	sterol	vegetable oil in bread	1.3	-12.6	3.9
Seki et al. 2003c #1	p	45	sterol	vegetable oil in bread	0.3	0.8	1.8
Seki et al. 2003c #2	p	44	sterol	vegetable oil in bread	0.5	-7.7	1.9
Seppe et al. 2007 study 1	p	60	stanol	low-fat yoghurt	2.0	-2.9	3.2
Seppe et al. 2007 study 2	p	61	stanol	low-fat yoghurt drink	2.0	-3.2	3.0
Seppe et al. 2007 study 3	p	19	stanol	low-fat yoghurt drink	2.0	-11.8	7.0
Seppe et al. 2007 study 4	p	59	stanol	low-fat milk	2.0	-6.2	2.4
Sialvera et al. 2011	p	108	sterol	yoghurt drink	4.0	-19.7	1.7
Sierksma et al. 1999	x	75	sterol	margarine	0.8	-6.1	0.6
Simons 2002 study 1	p	77	sterol	margarine	2.0	-10.2	2.8
Simons 2002 study 2	p	75	sterol	margarine	2.0	-6.1	3.6
Soderholm et al. 2011	p	63	sterol	rye bread	2.0	-8.1	3.5
Spilburg et al. 2003	p	24	stanol	lemonade	1.9	-14.3	4.5
Takeshita et al. 2008	p	29	sterol	DAG-containing cooking oil	0.5	-6.0	4.4
Temme et al. 2002	x	42	sterol	low-fat margarine	2.1	-9.6	1.4
Theuwissen & Mensink 2007	x	40	stanol	cereal	1.5	-4.4	2.1
Theuwissen et al. 2009	p	28	stanol	margarine	2.5	-9.5	4.6
Thomsen et al. 2004 #1	x	69	sterol	low-fat milk	1.2	-7.1	1.5
Thomsen et al. 2004 #2	x	69	sterol	low-fat milk	1.6	-9.6	1.5
Vanhanan 1994	p	14	stanol	mayonnaise	1.5	-2.0	7.6
Vanhanan et al. 1994	p	15	stanol	mayonnaise	0.8	-7.7	3.8
Vanstone et al. 2002 #1	x	15	sterol	butter	1.8	-10.2	2.8
Vanstone et al. 2002 #2	x	15	stanol	butter	1.8	-10.5	2.8
Vanstone et al. 2002 #3	x	15	mix	butter	1.8	-11.5	2.7
Varady et al. 2004 study 1	p	38	sterol	low-fat margarine	1.8	-11.3	2.4
Varady et al. 2004 study 2	p	36	sterol	low-fat margarine	1.8	-12.8	3.8
Volpe et al. 2001	x	30	sterol	low-fat yoghurt drink	1.1	-7.6	1.9
Weidner et al. 2008	p	50	sterol	soy drink	1.6	-5.2	2.8
Weststrate & Meijer 1998 #1	x	76	sterol	margarine	3.2	-13.1	0.6
Weststrate & Meijer 1998 #2	x	77	stanol	margarine	2.7	-11.9	0.6
Woodgate et al. 2006	p	29	stanol	softgel capsules	1.6	-7.2	3.7
Yoshida et al. 2006 study 1	x	16	mix	cereal bar	1.8	-6.1	2.5
Yoshida et al. 2006 study 2	x	13	mix	cereal bar	1.8	-2.8	3.7

¹Dose given as free equivalents in g/d, rounded off at 1 decimal.

² Multiple study arms in 1 study corrected for the same single control group/period: indicated with # 1, # 2, # 3, etc.

For some crossover studies, different active treatments were compared with a separate corresponding placebo treatment; however, as the same subjects were included in those periods, these are indicated with #1, #2, #3 etc.

³ Multiple study arms in 1 study, each corrected for a respective control group (i.e. different set of subjects): indicated with study 1, study 2, study 3, etc.

References

- AbuMweis SS, Vanstone CA, Ebine N, Kassis A, Ausman LM, Jones PJ, et al. (2006). Intake of a single morning dose of standard and novel plant sterol preparations for 4 weeks does not dramatically affect plasma lipid concentrations in humans. *The Journal of Nutrition* 136(4): 1012-1016.
- Algorta Pineda J, Chinchetrico Ranedo, MJ, Aguirre Anda, J, Francisco Terreros, S. (2005). Eficacia hipocolesterolimante de un yogur que contiene esteres de estanol vegetal. *Revista Clinica Espanola* 205(2): 63-66.
- Alhassan S, Reese KA, Mahurin J, Plaisance EP, Hilson BD, Garner JC, et al. (2006). Blood lipid responses to plant stanol ester supplementation and aerobic exercise training. *Metabolism: Clinical and Experimental* 55(4): 541-549.
- Andersson A, Karlström, B, Mohsen, R, Vessby, B. (1999). Cholesterol-lowering effects of a stanol ester-containing low-fat margarine used in conjunction with a strict lipid-lowering diet. *European Heart Journal Suppl* 1: S80-S90.
- Athyros VG, Kakafika AI, Papageorgiou AA, Tziomalos K, Peletidou A, Vosikis C, et al. (2011). Effect of a plant stanol ester-containing spread, placebo spread, or Mediterranean diet on estimated cardiovascular risk and lipid, inflammatory and haemostatic factors. *Nutrition, metabolism, and cardiovascular diseases : Nutrition, Metabolism and Cardiovascular Diseases* 21(3): 213-221.
- Banuls C, Martinez-Triguero ML, Lopez-Ruiz A, Morillas C, Jarabo MM, Bellod L, et al. (2011). Serum lipid responses to phytosterol-enriched milk in a moderate hypercholesterolemic population is not affected by apolipoprotein E polymorphism or diameter of low-density lipoprotein particles. *European Journal of Clinical Nutrition* 65(2): 255-261.
- Banuls C, Martinez-Triguero ML, Lopez-Ruiz A, Morillas C, Lacomba R, Victor VM, et al. (2010). Evaluation of cardiovascular risk and oxidative stress parameters in hypercholesterolemic subjects on a standard healthy diet including low-fat milk enriched with plant sterols. *Journal of Nutritional Biochemistry* 21(9): 881-886.
- Beer MU, Pritchard, H., Belsey, E.M., Davidson, M. (2000). Effect of a milk drink enriched with increasing doses of free tall oil phytosterols on plasma lipid levels of mildly hypercholesterolemic subjects. Document submitted to FDA. Docket No OOP1275 and OOP1276.
- Blair SN, Capuzzi DM, Gottlieb SO, Nguyen T, Morgan JM, Cater NB (2000). Incremental reduction of serum total cholesterol and low-density lipoprotein cholesterol with the addition of plant stanol ester-containing spread to statin therapy. *American Journal of Cardiology* 86(1): 46-52.
- Blomqvist S, Jauhainen, M, van Tol, A, Hyvönen, M, Torstila, M, Vanhanen, HT, Miettinen, TA, Ehnholm, C. (1993). Effect of sitostanol ester on composition and size distribution of low- and high-density lipoprotein. *Nutrition, Metabolism and Cardiovascular Diseases* 3: 158-164.
- Cater NB, Garcia-Garcia AB, Vega GL, Grundy SM (2005). Responsiveness of plasma lipids and lipoproteins to plant stanol esters. *American Journal of Cardiology* 96(1A): 23D-28D.
- Chen SC, Judd JT, Kramer M, Meijer GW, Clevidence BA, Baer DJ (2009). Phytosterol intake and dietary fat reduction are independent and additive in their ability to reduce plasma LDL cholesterol. *Lipids* 44(3): 273-281.
- Christiansen LI, Lahteenmaki PL, Mannelin MR, Seppanen-Laakso TE, Hiltunen RV, Yliruusi JK (2001). Cholesterol-lowering effect of spreads enriched with microcrystalline plant sterols in hypercholesterolemic subjects. *European Journal of Nutrition* 40(2): 66-73.
- Cleghorn CL, Skeaff CM, Mann J, Chisholm A (2003). Plant sterol-enriched spread enhances the cholesterol-lowering potential of a fat-reduced diet. *European Journal of Clinical Nutrition* 57(1): 170-176.
- Clifton PM, Mano M, Duchateau GS, van der Knaap HC, Trautwein EA (2008). Dose-response effects of different plant sterol sources in fat spreads on serum lipids and C-reactive protein and on the kinetic behavior of serum plant sterols. *European Journal of Clinical Nutrition* 62(8): 968-977.
- Clifton PM, Noakes M, Sullivan D, Erichsen N, Ross D, Annison G, et al. (2004). Cholesterol-lowering effects of plant sterol esters differ in milk, yoghurt, bread and cereal. *European Journal of Clinical Nutrition* 58(3): 503-509.
- Colgan HA, Floyd S, Noone EJ, Gibney MJ, Roche HM (2004). Increased intake of fruit and vegetables and a low-fat diet, with and without low-fat plant sterol-enriched spread consumption: effects on plasma lipoprotein and carotenoid metabolism. *Journal of Human Nutrition and Dietetics* 17(6): 561-569; quiz 571-564.
- Davidson MH, Maki KC, Umporowicz DM, Ingram KA, Dicklin MR, Schaefer E, et al. (2001). Safety and tolerability of esterified phytosterols administered in reduced-fat spread and salad dressing to healthy adult men and women. *Journal of the American College of Nutrition* 20(4): 307-319.
- De Graaf J, De Sauvage Nolting PR, Van Dam M, Belsey EM, Kastelein JJ, Haydn Pritchard P, et al. (2002). Consumption of tall oil-derived phytosterols in a chocolate matrix significantly decreases plasma total and low-density lipoprotein-cholesterol levels. *British Journal of Nutrition* 88(5): 479-488.
- De Jong A, Plat J, Bast A, Godschalk RW, Basu S, Mensink RP (2008a). Effects of plant sterol and stanol ester consumption on lipid metabolism, antioxidant status and markers of oxidative stress, endothelial function and low-grade inflammation in patients on current statin treatment. *European Journal of Clinical Nutrition* 62(2): 263-273.

- de Jong A, Plat J, Lutjohann D, Mensink RP (2008b). Effects of long-term plant sterol or stanol ester consumption on lipid and lipoprotein metabolism in subjects on statin treatment. *British Journal of Nutrition* 100(5): 937-941.
- Devaraj S, Autret BC, Jialal I (2006). Reduced-calorie orange juice beverage with plant sterols lowers C-reactive protein concentrations and improves the lipid profile in human volunteers. *American Journal of Clinical Nutrition* 84(4): 756-761.
- Devaraj S, Jialal I, Vega-Lopez S (2004). Plant sterol-fortified orange juice effectively lowers cholesterol levels in mildly hypercholesterolemic healthy individuals. *Arteriosclerosis Thrombosis and Vascular Biology* 24(3): e25-28.
- Doornbos AM, Meynen EM, Duchateau GS, van der Knaap HC, Trautwein EA (2006). Intake occasion affects the serum cholesterol lowering of a plant sterol-enriched single-dose yoghurt drink in mildly hypercholesterolaemic subjects. *European Journal of Clinical Nutrition* 60(3): 325-333.
- Eady S, Wallace A, Willis J, Scott R, Frampton C (2011). Consumption of a plant sterol-based spread derived from rice bran oil is effective at reducing plasma lipid levels in mildly hypercholesterolaemic individuals. *British Journal of Nutrition*: 1-12.
- Earnest CP, Mikus CR, Lemieux I, Arsenault BJ, Church TS (2007). Examination of encapsulated phytosterol ester supplementation on lipid indices associated with cardiovascular disease. *Nutrition* 23(9): 625-633.
- Escurriol V, Cofan, M, Ros, E. (2010). Fitoesteroles circulantes: biomarcadores de la absorcion de los esteroles de la dieta habitual y de adherencia a alimentos suplementados con esteroles vegetales. *Actividad Dietetica* 14(1): 3-9.
- Fuentes F, Lopez-Miranda J, Garcia A, Perez-Martinez P, Moreno J, Cofan M, et al. (2008). Basal plasma concentrations of plant sterols can predict LDL-C response to sitosterol in patients with familial hypercholesterolemia. *European Journal of Clinical Nutrition* 62(4): 495-501.
- Geelen A, Zock PL, de Vries JH, Katan MB (2002). Apolipoprotein E polymorphism and serum lipid response to plant sterols in humans. *European Journal of Clinical Investigation* 32(10): 738-742.
- Goldberg AC, Ostlund RE, Jr., Bateman JH, Schimmoeller L, McPherson TB, Spilburg CA (2006). Effect of plant stanol tablets on low-density lipoprotein cholesterol lowering in patients on statin drugs. *American Journal of Cardiology* 97(3): 376-379.
- Goncalves S, Vasco Maria, A, Silva-Herdade, AS, Martins e Silva, J, Saldanha, C. (2007). Milk enriched with phytosterols reduces plasma cholesterol levels in healthy and hypercholesterolemic subjects. *Nutrition Research* 27: 200-205.
- Gylling H, Radhakrishnan R, Miettinen T (1997). Reduction of serum cholesterol of postmenopausal women with previous myocardial infarction and cholesterol malabsorption induced by dietary sitostanol ester margarine: women and dietary sitostanol. *Circulation* 96(12): 4226-4231.
- Gylling H, Miettinen TA (1999). Cholesterol reduction by different plant stanol mixtures and with variable fat intake. *Metabolism: Clinical and Experimental* 48(5): 575-580.
- Gylling H, Miettinen TA (1996). Effects of inhibiting cholesterol absorption and synthesis on cholesterol and lipoprotein metabolism in hypercholesterolemic non-insulin-dependent diabetic men. *Journal of Lipid Research* 37(8): 1776-1785.
- Gylling H, Miettinen TA (1994). Serum cholesterol and cholesterol and lipoprotein metabolism in hypercholesterolaemic NIDDM patients before and during sitostanol ester-margarine treatment. *Diabetologia* 37(8): 773-780.
- Gylling H, Hallikainen M, Nissinen M, Miettinen TA (2010). The effect of a very high daily plant stanol ester intake on serum lipids, carotenoids, and fat-soluble vitamins. *Clinical Nutrition* 29: 112-118.
- Hallikainen M, Kurl S, Laakso M, Miettinen TA, Gylling H (2011). Plant stanol esters lower LDL cholesterol level in statin-treated subjects with type 1 diabetes by interfering the absorption and synthesis of cholesterol. *Atherosclerosis* 217(2): 473-478.
- Hallikainen M, Lyyra-Laitinen T, Laitinen T, Moilanen L, Miettinen TA, Gylling H (2008). Effects of plant stanol esters on serum cholesterol concentrations, relative markers of cholesterol metabolism and endothelial function in type 1 diabetes. *Atherosclerosis* 199(2): 432-439.
- Hallikainen MA, Sarkkinen ES, Gylling H, Erkkila AT, Uusitupa MI (2000). Comparison of the effects of plant sterol ester and plant stanol ester-enriched margarines in lowering serum cholesterol concentrations in hypercholesterolaemic subjects on a low-fat diet. *European Journal of Clinical Nutrition* 54(9): 715-725.
- Hallikainen MA, Uusitupa MI (1999). Effects of 2 low-fat stanol ester-containing margarines on serum cholesterol concentrations as part of a low-fat diet in hypercholesterolemic subjects. *American Journal of Clinical Nutrition* 69(3): 403-410.
- Hansel B, Nicolle C, Lalanne F, Tondu F, Lassel T, Donazzolo Y, et al. (2007). Effect of low-fat, fermented milk enriched with plant sterols on serum lipid profile and oxidative stress in moderate hypercholesterolemia. *American Journal of Clinical Nutrition* 86(3): 790-796.
- Hayes KC, Pronczuk A, Perlman D (2004). Nonesterified phytosterols dissolved and recrystallized in oil reduce plasma cholesterol in gerbils and humans. *The Journal of Nutrition* 134(6): 1395-1399.
- Heggen E, Granlund L, Pedersen JI, Holme I, Ceglarek U, Thiery J, et al. (2010). Plant sterols from rapeseed and tall oils: effects on lipids, fat-soluble vitamins and plant sterol concentrations. *Nutrition, Metabolism and Cardiovascular Diseases* 20(4): 258-265.
- Hendriks HF, Brink EJ, Meijer GW, Princen HM, Ntanios FY (2003). Safety of long-term consumption of plant sterol esters-enriched spread. *European Journal of Clinical Nutrition* 57(5): 681-692.

Hendriks HF, Weststrate JA, van Vliet T, Meijer GW (1999). Spreads enriched with three different levels of vegetable oil sterols and the degree of cholesterol lowering in normocholesterolaemic and mildly hypercholesterolaemic subjects. European Journal of Clinical Nutrition 53(4): 319-327.

Hernandez-Mijares A, Banuls C, Jover A, Sola E, Bellod L, Martinez-Triguero ML, et al. (2011). Low intestinal cholesterol absorption is associated with a reduced efficacy of phytosterol esters as hypolipemic agents in patients with metabolic syndrome. Clinical Nutrition 30(5): 604-609.

Hernandez-Mijares A, Banuls C, Rocha M, Morillas C, Martinez-Triguero ML, Victor VM, et al. (2010). Effects of phytosterol ester-enriched low-fat milk on serum lipoprotein profile in mildly hypercholesterolaemic patients are not related to dietary cholesterol or saturated fat intake. British Journal of Nutrition 104(7): 1018-1025.

Hironaka T, Shioya, N., Matsubara, H., Matsuoka, Y., Itakura, H. (2006). Double-blind, placebo-controlled study of effects of plant sterol enriched vegetable juice on serum cholesterol concentrations in mildly hypercholesterolemic subjects and safety evaluation. Journal of Oleo Science. 55(11): 593-606.

Homma Y, Ikeda I, Ishikawa T, Tateno M, Sugano M, Nakamura H (2003). Decrease in plasma low-density lipoprotein cholesterol, apolipoprotein B, cholestryler ester transfer protein, and oxidized low-density lipoprotein by plant stanol ester-containing spread: a randomized, placebo-controlled trial. Nutrition 19(4): 369-374.

Houweling AH, Vanstone CA, Trautwein EA, Duchateau GS, Jones PJ (2009). Baseline plasma plant sterol concentrations do not predict changes in serum lipids, C-reactive protein (CRP) and plasma plant sterols following intake of a plant sterol-enriched food. European Journal of Clinical Nutrition 63(4): 543-551.

Hyun Y, Kim OY, Kang, JB, Lee, JH, Jang, Y, Liponkoski, L, Salo, P. (2005). Plant stanol esters in low-fat yogurt reduces total and low-density lipoprotein cholesterol and low-density lipoprotein oxidation in normocholesterolemic and mildly hypercholesterolemic subjects. Nutrition Research 25: 743-753.

Ishizaki T, Wakabayashi, M, Tanimoto, H, Shima, A, Yabune, M, Kajimoto, O, Itakura, H. (2003). Effects of long-term intake of mayonnaise containing phytosterolester on blood cholesterol concentration in japanese with borderline or mild cholesterolemia. Journal of Clinical Biochemistry and Nutrition 33(3): 75-82

Jakulj L, Trip MD, Sudhop T, von Bergmann K, Kastelein JJ, Vissers MN (2005). Inhibition of cholesterol absorption by the combination of dietary plant sterols and ezetimibe: effects on plasma lipid levels. Journal of Lipid Research 46(12): 2692-2698.

Jauhainen T, Salo P, Niittynen L, Poussa T, Korppela R (2006). Effects of low-fat hard cheese enriched with plant stanol esters on serum lipids and apolipoprotein B in mildly hypercholesterolaemic subjects. European Journal of Clinical Nutrition 60(11): 1253-1257.

Jones PJ, Ntanios FY, Raeini-Sarjaz M, Vanstone CA (1999). Cholesterol-lowering efficacy of a sitostanol-containing phytosterol mixture with a prudent diet in hyperlipidemic men. American Journal of Clinical Nutrition 69(6): 1144-1150.

Jones PJ, Raeini-Sarjaz M, Ntanios FY, Vanstone CA, Feng JY, Parsons WE (2000). Modulation of plasma lipid levels and cholesterol kinetics by phytosterol versus phytostanol esters. Journal of Lipid Research 41(5): 697-705.

Jones PJ, Vanstone CA, Raeini-Sarjaz M, St-Onge MP (2003). Phytosterols in low- and nonfat beverages as part of a controlled diet fail to lower plasma lipid levels. Journal of Lipid Research 44(9): 1713-1719.

Judd JT, Baer DJ, Chen SC, Clevidence BA, Muesing RA, Kramer M, et al. (2002). Plant sterol esters lower plasma lipids and most carotenoids in mildly hypercholesterolemic adults. Lipids 37(1): 33-42.

Kassis AN, Vanstone CA, AbuMweis SS, Jones PJ (2008). Efficacy of plant sterols is not influenced by dietary cholesterol intake in hypercholesterolemic individuals. Metabolism: Clinical and Experimental 57(3): 339-346.

Khandelwal S, Demonty I, Jeemon P, Lakshmy R, Mukherjee R, Gupta R, et al. (2009). Independent and interactive effects of plant sterols and fish oil n-3 long-chain polyunsaturated fatty acids on the plasma lipid profile of mildly hyperlipidaemic Indian adults. British Journal of Nutrition 102(5): 722-732.

Korpela R, Tuomilehto J, Hogstrom P, Seppo L, Piironen V, Salo-Vaananten P, et al. (2006). Safety aspects and cholesterol-lowering efficacy of low fat dairy products containing plant sterols. European Journal of Clinical Nutrition 60(5): 633-642.

Kratz M, Kannenberg F, Gramenz E, Berning B, Trautwein E, Assmann G, et al. (2007). Similar serum plant sterol responses of human subjects heterozygous for a mutation causing sitosterolemia and controls to diets enriched in plant sterols or stanols. European Journal of Clinical Nutrition 61(7): 896-905.

Kurokawa M, Masuda Y, Noda M, Marushima R, Usuda M, Takeda S, et al. (2008). [Minimal effective dose on serum cholesterol concentration and the safety evaluation of dressing containing plant sterol in Japanese subjects]. Journal of Oleo Science 57(1): 23-33.

Kurokawa M, Masuda, Y, Noda, M, Usuda, M, Takeda, S, Hasegawa, M, Homma, Y, Sugano, M. (2008). Effects of dressing containing plant sterol on serum cholesterol concentration and the safety evaluation in borderline or mildly hypercholesterolemic Japanese subjects. Journal of Oleo Science 57(1): 35-45.

Lagstrom H, Helenius, H, Salo, P. (2007). Serum cholesterol-lowering efficacy of stanol ester incorporated in gelatin capsules. Scandinavian Journal of Food and Nutrition 50(3): 124-130.

- Lau VW, Journoud M, Jones PJ (2005). Plant sterols are efficacious in lowering plasma LDL and non-HDL cholesterol in hypercholesterolemic type 2 diabetic and nondiabetic persons. *American Journal of Clinical Nutrition* 81(6): 1351-1358.
- Lee YM, Haastert B, Scherbaum W, Hauner H (2003). A phytosterol-enriched spread improves the lipid profile of subjects with type 2 diabetes mellitus--a randomized controlled trial under free-living conditions. *European Journal of Nutrition* 42(2): 111-117.
- Li NY, Li K, Qi Z, Demonty I, Gordon M, Francis L, et al. (2007). Plant sterol-enriched milk tea decreases blood cholesterol concentrations in Chinese adults: a randomised controlled trial. *British Journal of Nutrition* 98(5): 978-983.
- Lin X, Racette SB, Lefevre M, Ma L, Spearie CA, Steger-May K, et al. (2011). Combined effects of ezetimibe and phytosterols on cholesterol metabolism: a randomized, controlled feeding study in humans. *Circulation* 124(5): 596-601.
- Lottenberg AM, Nunes VS, Nakandakare ER, Neves M, Bernik M, Lagrost L, et al. (2003). The human cholesteryl ester transfer protein I405V polymorphism is associated with plasma cholesterol concentration and its reduction by dietary phytosterol esters. *The Journal of Nutrition* 133(6): 1800-1805.
- Madsen MB, Jensen AM, Schmidt EB (2007). The effect of a combination of plant sterol-enriched foods in mildly hypercholesterolemic subjects. *Clinical Nutrition* 26(6): 792-798.
- Maki KC, Davidson MH, Umporowicz DM, Schaefer EJ, Dicklin MR, Ingram KA, et al. (2001). Lipid responses to plant-sterol-enriched reduced-fat spreads incorporated into a National Cholesterol Education Program Step I diet. *American Journal of Clinical Nutrition* 74(1): 33-43.
- Mannarino E, Pirro M, Cortese C, Lupattelli G, Siepi D, Mezzetti A, et al. (2009). Effects of a phytosterol-enriched dairy product on lipids, sterols and 8-isoprostane in hypercholesterolemic patients: a multicenter Italian study. *Nutrition, Metabolism and Cardiovascular Diseases*. 19(2): 84-90.
- Matsuoka R, Masuda, Y., Takeuchi, A., Marushima, R., Hasegawa, M, Sakamoto, A, Hirata, H, Kajimoto, O, Homma, Y. (2004a). A double-blind, placebo-controlled study on the effects of mayonnaise containing free plant sterol on serum cholesterol concentration; safety evaluation for normocholesterolemic and mildly hypercholesterolemic Japanese subjects. *Journal of Oleo Science* 53(2): 79-88.
- Matsuoka R, Masuda, Y., Takeuchi, A., Marushima, R., Onuki, M. (2004b). Minimal effective dose of plant sterol on serum cholesterol concentration in Japanese subjects and safety evaluation of plant sterol supplemented in mayonnaise. *Journal of Oleo Science* 53(1): 17-27.
- Matvienko OA, Lewis DS, Swanson M, Arndt B, Rainwater DL, Stewart J, et al. (2002). A single daily dose of soybean phytosterols in ground beef decreases serum total cholesterol and LDL cholesterol in young, mildly hypercholesterolemic men. *American Journal of Clinical Nutrition* 76(1): 57-64.
- McPherson TB, Ostlund RE, Goldberg AC, Bateman JH, Schimmoeller L, Spilburg CA (2005). Phytostanol tablets reduce human LDL-cholesterol. *Journal of Pharmacy and Pharmacology* 57(7): 889-896.
- Mensink RP, de Jong A, Lutjohann D, Haenen GR, Plat J (2010). Plant stanols dose-dependently decrease LDL-cholesterol concentrations, but not cholesterol-standardized fat-soluble antioxidant concentrations, at intakes up to 9 g/d. *American Journal of Clinical Nutrition* 92(1): 24-33.
- Mensink RP, Ebbing S, Lindhout M, Plat J, van Heugten MM (2002). Effects of plant stanol esters supplied in low-fat yoghurt on serum lipids and lipoproteins, non-cholesterol sterols and fat soluble antioxidant concentrations. *Atherosclerosis* 160(1): 205-213.
- Miettinen TA, Puska P, Gylling H, Vanhanen H, Vartiainen E (1995). Reduction of serum cholesterol with sitostanol-ester margarine in a mildly hypercholesterolemic population. *New England Journal of Medicine* 333(20): 1308-1312.
- Miettinen TA, Vanhanen H (1994). Dietary sitostanol related to absorption, synthesis and serum level of cholesterol in different apolipoprotein E phenotypes. *Atherosclerosis* 105(2): 217-226.
- Mussner MJ, Parhofer KG, Von Bergmann K, Schwandt P, Broedl U, Otto C (2002). Effects of phytosterol ester-enriched margarine on plasma lipoproteins in mild to moderate hypercholesterolemia are related to basal cholesterol and fat intake. *Metabolism* 51(2): 189-194.
- Naumann E, Plat J, Mensink RP (2003). Changes in serum concentrations of noncholesterol sterols and lipoproteins in healthy subjects do not depend on the ratio of plant sterols to stanols in the diet. *The Journal of Nutrition* 133(9): 2741-2747.
- Neil HA, Meijer GW, Roe LS (2001). Randomised controlled trial of use by hypercholesterolaemic patients of a vegetable oil sterol-enriched fat spread. *Atherosclerosis* 156(2): 329-337.
- Nestel P, Cehun M, Pomeroy S, Abbey M, Weldon G (2001). Cholesterol-lowering effects of plant sterol esters and non-esterified stanols in margarine, butter and low-fat foods. *European Journal of Clinical Nutrition* 55(12): 1084-1090.
- Nigon F, Serfaty-Lacroix C, Beucler I, Chauvois D, Neveu C, Giral P, et al. (2001). Plant sterol-enriched margarine lowers plasma LDL in hyperlipidemic subjects with low cholesterol intake: effect of fibrate treatment. *Clinical Chemistry and Laboratory Medicine* 39(7): 634-640.
- Niittynen L, Jauhainen, TA, Poussa, TA, Korpela, R. (2007). Effects of yoghurt enriched with free plant sterols on the levels of serum lipids and plant sterols in moderately hypercholesterolaemic subjects on a high-fat diet. *International Journal of Food Sciences and Nutrition* 1: 1-11.

- Noakes M, Clifton P, Ntanios F, Shrapnel W, Record I, McInerney J (2002). An increase in dietary carotenoids when consuming plant sterols or stanols is effective in maintaining plasma carotenoid concentrations. *American Journal of Clinical Nutrition* 75(1): 79-86.
- Noakes M, Clifton PM, Doornbos AM, Trautwein EA (2005). Plant sterol ester-enriched milk and yoghurt effectively reduce serum cholesterol in modestly hypercholesterolemic subjects. *European Journal of Nutrition* 44(4): 214-222.
- Ntanios FY, Homma Y, Ushiro S (2002). A spread enriched with plant sterol-esters lowers blood cholesterol and lipoproteins without affecting vitamins A and E in normal and hypercholesterolemic Japanese men and women. *Journal of Nutrition* 132(12): 3650-3655.
- Ooi EM, Watts GF, Barrett PH, Chan DC, Clifton PM, Ji J, et al. (2007). Dietary plant sterols supplementation does not alter lipoprotein kinetics in men with the metabolic syndrome. *Asia Pacific Journal of Clinical Nutrition* 16(4): 624-631.
- Pelletier X, Belbraouet S, Mirabel D, Mordret F, Perrin JL, Pages X, et al. (1995). A diet moderately enriched in phytosterols lowers plasma cholesterol concentrations in normocholesterolemic humans. *Annals of Nutrition and Metabolism* 39(5): 291-295.
- Plana N, Nicolle C, Ferre R, Camps J, Cos R, Villoria J, et al. (2008). Plant sterol-enriched fermented milk enhances the attainment of LDL-cholesterol goal in hypercholesterolemic subjects. *European Journal of Nutrition* 47(1): 32-39.
- Plat J, Mensink RP (2000). Vegetable oil based versus wood based stanol ester mixtures: effects on serum lipids and hemostatic factors in non-hypercholesterolemic subjects. *Atherosclerosis* 148(1): 101-112.
- Plat J, van Onselen EN, van Heugten MM, Mensink RP (2000). Effects on serum lipids, lipoproteins and fat soluble antioxidant concentrations of consumption frequency of margarines and shortenings enriched with plant stanol esters. *European Journal of Clinical Nutrition* 54(9): 671-677.
- Quilez J, Rafecas M, Brufau G, Garcia-Lorda P, Megias I, Bullo M, et al. (2003). Bakery products enriched with phytosterol esters, alpha-tocopherol and beta-carotene decrease plasma LDL-cholesterol and maintain plasma beta-carotene concentrations in normocholesterolemic men and women. *Journal of Nutrition* 133(10): 3103-3109.
- Racette SB, Lin X, Lefevre M, Spearie CA, Most MM, Ma L, et al. (2010). Dose effects of dietary phytosterols on cholesterol metabolism: a controlled feeding study. *American Journal of Clinical Nutrition* 91(1): 32-38.
- Raitakari OT, Salo P, Gylling H, Miettinen TA (2008). Plant stanol ester consumption and arterial elasticity and endothelial function. *British Journal of Nutrition* 100(3): 603-608.
- Rudkowska I, AbuMweis SS, Nicolle C, Jones PJ (2008). Cholesterol-lowering efficacy of plant sterols in low-fat yogurt consumed as a snack or with a meal. *Journal of the American College of Nutrition* 27(5): 588-595.
- Ruiu G, Pinach S, Veglia F, Gambino R, Marena S, Uberti B, et al. (2009). Phytosterol-enriched yogurt increases LDL affinity and reduces CD36 expression in polygenic hypercholesterolemia. *Lipids* 44(2): 153-160.
- Saito S, Takeshita M, Tomonobu K, Kudo N, Shiiba D, Hase T, et al. (2006). Dose-dependent cholesterol-lowering effect of a mayonnaise-type product with a main component of diacylglycerol-containing plant sterol esters. *Nutrition* 22(2): 174-178.
- Seki S, Hidaka I, Kojima K, Yoshino H, Aoyama T, Okazaki M, et al. (2003a). Effects of phytosterol ester-enriched vegetable oil on plasma lipoproteins in healthy men. *Asia Pacific Journal of Clinical Nutrition* 12(3): 282-291.
- Seki S, Abe, T., Hidaka, I., Kojima, K., Yoshino, H., Aoyama, T., Okazaki, M., Kondo, K. (2003b). Effects of phytosterol ester-enriched vegetable oil on serum cholesterol and assessment of safety in healthy men. *Journal of Oleo Science* 52(4): 205-213.
- Seki S, Abe, T., Hidaka, I., Kojima, K., Yoshino, H., Aoyama, T., Okazaki, M., Kondo, K. (2003c). Vegetable oil enriched with phytosterol ester and the degree of serum cholesterol lowering effect in healthy subjects. *Journal of Oleo Science* 52(6): 285-294.
- Seppo L, Jauhainen T, Nevala R, Poussa T, Korpela R (2007). Plant stanol esters in low-fat milk products lower serum total and LDL cholesterol. *European Journal of Nutrition* 46(2): 111-117.
- Sialvera TE, Pounis GD, Koutelidakis AE, Richter DJ, Yfanti G, Kapsokafalou M, Goumas G, Chiotinis N, Diamantopoulos E, Zampelas A (2012). Phytosterol supplementation decreases plasma small and dense LDL levels in metabolic syndrome patients on a westernized type diet. *Nutrition, Metabolism and Cardiovascular Diseases* 22(10): 843-848.
- Siersma A, Weststrate JA, Meijer GW (1999). Spreads enriched with plant sterols, either esterified 4,4-dimethylsterols or free 4-desmethylsterols, and plasma total- and LDL-cholesterol concentrations. *British Journal of Nutrition* 82(4): 273-282.
- Simons LA (2002). Additive effect of plant sterol-ester margarine and cerivastatin in lowering low-density lipoprotein cholesterol in primary hypercholesterolemia. *The American Journal of Cardiology* 90(7): 737-740.
- Soderholm PP, Alftan G, Koskela AH, Adlercreutz H, Tikkannen MJ (2012). The effect of high-fiber rye bread enriched with nonesterified plant sterols on major serum lipids and apolipoproteins in normocholesterolemic individuals. *Nutrition, Metabolism and Cardiovascular Diseases* 22(7): 575-582.
- Spilburg CA, Goldberg AC, McGill JB, Stenson WF, Racette SB, Bateman J, et al. (2003). Fat-free foods supplemented with soy stanol-lecithin powder reduce cholesterol absorption and LDL cholesterol. *Journal of the American Dietetic Association* 103(5): 577-581.
- Takeshita M, Katsuragi, Y., Kusuhara, M, Higashi, K., Miyajima, E, Mizuno, K, Mori, K, Obata, T., Ohmori, R., Ohsuzu, F., Onodera, Y., Sano, J., Sawada, S., Tabata, S., Tokimitsu, I., Tomonobu, K., Yamashita, T., Yasukawa, T., Yonemura, A., Nakamura, H.

- (2008). Phytosterols dissolved in diacylglycerol oil reinforce the cholesterol-lowering effect of low-dose pravastatin treatment. *Nutrition, Metabolism and Cardiovascular Diseases* 18: 483-491.
- Temme EH, Van Hoydonck PG, Schouten EG, Kesteloot H (2002). Effects of a plant sterol-enriched spread on serum lipids and lipoproteins in mildly hypercholesterolaemic subjects. *Acta Cardiologica* 57(2): 111-115.
- Theuwissen E, Mensink RP (2007). Simultaneous intake of beta-glucan and plant stanol esters affects lipid metabolism in slightly hypercholesterolemic subjects. *The Journal of Nutrition* 137(3): 583-588.
- Theuwissen E, Plat J, van der Kallen CJ, van Greevenbroek MM, Mensink RP (2009). Plant stanol supplementation decreases serum triacylglycerols in subjects with overt hypertriglyceridemia. *Lipids* 44(12): 1131-1140.
- Thomsen AB, Hansen HB, Christiansen C, Green H, Berger A (2004). Effect of free plant sterols in low-fat milk on serum lipid profile in hypercholesterolemic subjects. *European Journal of Clinical Nutrition* 58(6): 860-870.
- Vanhanan H (1994). Cholesterol malabsorption caused by sitostanol ester feeding and neomycin in pravastatin-treated hypercholesterolaemic patients. *European Journal of Clinical Pharmacology* 47(2): 169-176.
- Vanhanan HT, Kajander J, Lehtovirta H, Miettinen TA (1994). Serum levels, absorption efficiency, faecal elimination and synthesis of cholesterol during increasing doses of dietary sitostanol esters in hypercholesterolaemic subjects. *Clinical Science (Lond)* 87(1): 61-67.
- Vanstone C, Raini-Sarjaz, M, Parsons, WE, Jones, PJH. (2002). Unesterified plant sterols and stanols lower LDL-cholesterol concentrations equivalently in hypercholesterolemic persons. *American Journal of Clinical Nutrition* 76: 1272-1278.
- Varady KA, Ebine N, Vanstone CA, Parsons WE, Jones PJ (2004). Plant sterols and endurance training combine to favorably alter plasma lipid profiles in previously sedentary hypercholesterolemic adults after 8 wk. *American Journal of Clinical Nutrition* 80(5): 1159-1166.
- Volpe R, Niittynen L, Korpela R, Sirtori C, Bucci A, Fraone N, et al. (2001). Effects of yoghurt enriched with plant sterols on serum lipids in patients with moderate hypercholesterolemia. *British Journal of Nutrition* 86(2): 233-239.
- Weidner C, Krempf M, Bard JM, Cazaubiel M, Bell D (2008). Cholesterol lowering effect of a soy drink enriched with plant sterols in a French population with moderate hypercholesterolemia. *Lipids Health Dis* 7: 35.
- Weststrate JA, Meijer GW (1998). Plant sterol-enriched margarines and reduction of plasma total- and LDL-cholesterol concentrations in normocholesterolaemic and mildly hypercholesterolaemic subjects. *European Journal of Clinical Nutrition* 52(5): 334-343.
- Woodgate D, Chan CH, Conquer JA (2006). Cholesterol-lowering ability of a phytostanol softgel supplement in adults with mild to moderate hypercholesterolemia. *Lipids* 41(2): 127-132.
- Yoshida M, Vanstone CA, Parsons WD, Zawistowski J, Jones PJ (2006). Effect of plant sterols and glucomannan on lipids in individuals with and without type II diabetes. *European Journal of Clinical Nutrition* 60(4): 529-537.