

Prenatal vitamin D3 supplementation suppresses LL-37 peptide expression in ex vivo activated neonatal macrophages but not their killing capacity

Rubhana Raqib¹, Anna Ly², Evana Akhtar¹, Akhirunnesa Mily¹, Nandita Perumal², Abdullah Al-Mahmud¹, Birgitta Agerberth³, Abdullah Hel Baqui⁴, Daniel E. Roth²

¹Nutritional Biochemistry Laboratory, International Centre for Diarrheal Disease Research, Bangladesh, 1212 Dhaka, Bangladesh; ²Department of Paediatrics, Hospital for Sick Children and University of Toronto, 555 University Avenue, Toronto, ON M5G 1X8, Canada; ³Department of Medical Biochemistry and Biophysics, Karolinska Institutet, 171 77 Stockholm, Sweden; ⁴ International Center for Maternal and Newborn Health, Department of International Health, The Johns Hopkins Bloomberg School of Public Health, 615 North Wolfe Street, Baltimore, MD 21205, USA

Correspondence to:

Rubhana Raqib, Nutritional Biochemistry Laboratory, Centre for Vaccine Sciences, International Centre for Diarrheal Disease Research, Bangladesh (icddr,b), Mohakhali, Dhaka-1212, Bangladesh, Fax: +880-28812529; Phone- 880-2-9827068; Email: rubhana@icddrb.org

Supplementary Table 1. Concentration of 25(OH)D in maternal blood at baseline, delivery and in cord blood by treatment group.

Serum 25(OH)D	Treatment groups ¹	
	VitD (n=63)	Placebo (n=66)
Baseline	43 (30, 60)	41 (27, 56)
Delivery	133 (112, 159)	34 (28, 48)
Cord blood	100 (83, 119)	35 (27, 46)

¹ Median with Interquartile range in parentheses unless otherwise indicated. Concentrations of 25(OH)D are expressed in nmol/L.