


Supplemental Figure I. Participants' flow chart



Supplemental Table I. Participants' characteristics for included and excluded subjects ^a

	Included subjects	Excluded subjects	P*
Age, year	55.9 ± 10.3	55.8 ± 9.5	0.09
Males, %	36.9	39.9	<0.01
Body mass index, kg/m ²	22.7 ± 2.8	22.7 ± 2.8	0.63
Ethanol intake, g/day	28.9 ± 22.5	29.0 ± 23.0	0.51
Thiamine intake, mg/day	0.75 ± 0.23	0.75 ± 0.23	0.98
Riboflavin intake, mg/day	1.3 ± 0.4	1.3 ± 0.5	0.93
Niacin intake, mg/day	13.2 ± 4.2	13.2 ± 4.2	0.92
Pantothenic acid intake, mg/day	5.3 ± 1.5	5.3 ± 1.5	0.66
Pyridoxine intake, mg/day	1.1 ± 0.3	1.1 ± 0.3	0.32
Folate intake, mg/day	418.9 ± 161.3	419.1 ± 160.8	0.90
Cobalamin intake, mg/day	7.6 ± 0.4	7.6 ± 0.5	0.89
Ascorbic acid intake, mg/day	140.0 ± 49.9	139.4 ± 50.1	0.19
Family history of diabetes, %	3.0	2.9	0.51
History of hypertension, %	16.3	18.7	<0.01
Current smoker, %	23.9	24.2	0.44
Sports ≥5 hour/week, %	11.3	10.9	0.16
Walking ≥5 hour/week, %	54.3	53.5	0.08
≥ 1 cup of green tea/day, %	70.6	74.8	<0.01
≥ 1 cup of coffee/day, %	37.1	31.7	<0.01
Daily use of multivitamins, %	3.5	3.6	0.55
Daily use of vitamin B1, %	1.3	1.4	0.33
Daily use of vitamin C, %	3.6	3.5	0.08
Daily use of vitamin E, %	3.6	3.6	0.91

^a Excluded subjects are those with missing data for the history of diabetes at baseline or the 5-year follow-up.

*Student t-test (sometimes with log transformation whenever needed) was used for continuous variables and Chi Square test was used for categorical variables.