

## Data supplement

**Table DS1** Rates of mental health problems by cumulative history of abortion, pregnancy loss and live birth with or without an adverse reaction to pregnancy<sup>a</sup>

| Measure                               | Abortion         |      | Pregnancy loss   |      | Unwanted/adverse reaction |      | Live birth       |      |
|---------------------------------------|------------------|------|------------------|------|---------------------------|------|------------------|------|
|                                       | No               | Yes  | No               | Yes  | No                        | Yes  | No               | Yes  |
| Major depression, %                   |                  |      |                  |      |                           |      |                  |      |
| 15–18 years                           | 31.4             | 55.6 | 32.0             | 66.7 | 32.4                      | 50.0 | 32.5             | 38.5 |
| 18–21 years                           | 29.0             | 40.9 | 28.6             | 59.4 | 30.7                      | 26.9 | 30.7             | 29.1 |
| 21–25 years                           | 24.7             | 39.8 | 26.3             | 36.2 | 25.7                      | 45.5 | 26.1             | 32.4 |
| 25–30 years                           | 21.9             | 33.9 | 23.2             | 30.9 | 22.2                      | 47.1 | 25.0             | 24.1 |
| Pooled OR (95% CI)                    | 2.15 (1.44–3.23) |      | 1.94 (1.21–3.10) |      | 2.25 (1.28–3.94)          |      | 0.89 (0.60–1.30) |      |
| Anxiety disorder, %                   |                  |      |                  |      |                           |      |                  |      |
| 15–18 years                           | 38.0             | 51.9 | 38.5             | 55.6 | 38.8                      | 37.5 | 38.8             | 38.5 |
| 18–21 years                           | 17.2             | 22.7 | 16.4             | 40.6 | 18.0                      | 15.4 | 15.9             | 34.6 |
| 21–25 years                           | 20.4             | 38.7 | 21.7             | 39.7 | 22.6                      | 36.4 | 21.7             | 31.5 |
| 25–30 years                           | 21.4             | 33.0 | 22.0             | 33.0 | 22.4                      | 39.2 | 23.4             | 25.1 |
| Pooled OR (95% CI)                    | 2.25 (1.49–3.42) |      | 2.34 (1.45–3.77) |      | 1.85 (1.02–3.35)          |      | 1.34 (0.91–1.98) |      |
| Suicidal ideation, %                  |                  |      |                  |      |                           |      |                  |      |
| 15–18 years                           | 23.3             | 37.0 | 23.5             | 55.6 | 24.0                      | 25.0 | 24.3             | 15.4 |
| 18–21 years                           | 13.4             | 25.8 | 13.5             | 37.5 | 15.1                      | 11.5 | 14.1             | 21.8 |
| 21–25 years                           | 9.5              | 23.7 | 10.1             | 27.6 | 11.1                      | 22.7 | 10.6             | 17.6 |
| 25–30 years                           | 7.1              | 8.7  | 5.8              | 14.9 | 6.8                       | 13.7 | 6.1              | 9.7  |
| Pooled OR (95% CI)                    | 2.25 (1.30–3.89) |      | 3.30 (1.79–6.09) |      | 1.77 (0.80–3.94)          |      | 1.41 (0.83–2.39) |      |
| Alcohol dependence, %                 |                  |      |                  |      |                           |      |                  |      |
| 15–18 years                           | 5.5              | 0.0  | 5.1              | 11.1 | 5.3                       | 0.0  | 5.3              | 0.0  |
| 18–21 years                           | 4.7              | 4.6  | 3.5              | 21.9 | 4.9                       | 0.0  | 4.6              | 5.5  |
| 21–25 years                           | 2.6              | 7.5  | 3.1              | 6.9  | 3.6                       | 2.3  | 3.2              | 4.6  |
| 25–30 years                           | 1.8              | 5.2  | 2.4              | 3.2  | 2.4                       | 3.9  | 2.9              | 2.1  |
| Pooled OR (95% CI)                    | 1.86 (0.84–4.12) |      | 2.94 (1.24–6.96) |      | 0.54 (0.13–2.23)          |      | 1.05 (0.45–2.41) |      |
| Illicit drug dependence, %            |                  |      |                  |      |                           |      |                  |      |
| 15–18 years                           | 4.1              | 0.0  | 3.9              | 0.0  | 3.9                       | 0.0  | 3.9              | 0.0  |
| 18–21 years                           | 2.5              | 16.7 | 3.9              | 9.4  | 3.7                       | 15.4 | 3.9              | 7.3  |
| 21–25 years                           | 3.1              | 9.7  | 3.7              | 8.6  | 4.0                       | 6.8  | 4.2              | 4.6  |
| 25–30 years                           | 1.5              | 8.7  | 2.7              | 5.3  | 2.9                       | 5.9  | 3.2              | 3.1  |
| Pooled OR (95% CI)                    | 7.08 (3.06–16.4) |      | 1.76 (0.58–5.35) |      | 2.11 (0.68–6.54)          |      | 0.72 (0.28–1.86) |      |
| Mean number of mental health problems |                  |      |                  |      |                           |      |                  |      |
| 15–18 years                           | 1.02             | 1.44 | 1.03             | 1.89 | 1.04                      | 1.13 | 1.05             | 0.92 |
| 18–21 years                           | 0.67             | 1.11 | 0.66             | 1.69 | 0.72                      | 0.69 | 0.69             | 0.98 |
| 21–25 years                           | 0.60             | 1.19 | 0.65             | 1.19 | 0.67                      | 1.14 | 0.66             | 0.91 |
| 25–30 years                           | 0.54             | 0.90 | 0.56             | 0.87 | 0.57                      | 1.10 | 0.61             | 0.64 |
| Pooled IRR (95% CI)                   | 1.54 (1.28–1.85) |      | 1.49 (1.21–1.84) |      | 1.31 (1.01–1.69)          |      | 1.03 (0.86–1.22) |      |
| Sample sizes, n                       |                  |      |                  |      |                           |      |                  |      |
| 15–18 years                           | 497              | 27   | 515              | 9    | 516                       | 8    | 511              | 13   |
| 18–21 years                           | 449              | 66   | 483              | 32   | 489                       | 26   | 460              | 55   |
| 21–25 years                           | 421              | 93   | 456              | 58   | 470                       | 44   | 406              | 108  |
| 25–30 years                           | 392              | 115  | 413              | 94   | 456                       | 51   | 312              | 195  |

IRR, incident rate ratio.

a. Assessed up to the end of each interval.