

Data supplement

| Table DS1 Demographic characteristics of the total sample across groups | | | | |
|--|---------------------------|-----------------------------------|-----------------------------------|---------------------------------|
| | Total (<i>n</i> = 56) | Inplant group (<i>n</i> = 29) | Control group (<i>n</i> = 27) | <i>P</i> value of difference |
| Female, <i>n</i> (%) | 20 (36) | 11 (38) | 9 (33) | 0.79 |
| Age, years: mean (s.d.) | 34.2 (8.6) | 34.5 (8) | 34 (9.4) | 0.84 |
| Age 18–30 years, <i>n</i> (%) | 24 (43) | 11 (35) | 13 (48) | |
| Age 31–59 years, <i>n</i> (%) | 32 (57) | 18 (65) | 14 (52) | |
| Age range, years | 21–55 | 22–50 | 21–55 | |
| Immigrant (first or second generation), <i>n</i> (%) | 9 (16) | 5 (17) | 4 (15) | 0.55 |
| Pre-inclusion length of stay, weeks: mean (s.d.) | 20 (38) | 27 (34.2) | 12.6 (42) | 0.17 |
| Lifetime substance use, years: mean (s.d.) | | | | |
| Heroin | 7 (5.3) | 7 (5) | 6.9 (5.6) | 0.98 |
| Methadone | 1 (1.7) | 1.1 (1.8) | 0.8 (1.5) | 0.46 |
| Other opioids | 2.2 (4.2) | 2.8 (5.1) | 1.6 (3.1) | 0.30 |
| Benzodiazepines | 5.4 (6.4) | 7.3 (7.2) | 3.3 (4.8) | 0.02* |
| Heavy alcohol use | 3.3 (5.9) | 3.9 (7.1) | 2.8 (4.3) | 0.56 |
| Amphetamine | 4.3 (5.2) | 4.7 (4.9) | 4 (5.6) | 0.62 |
| Cocaine | 0.5 (1.3) | 0.4 (1.1) | 0.6 (1.5) | 0.68 |
| Cannabis | 10.1 (7.7) | 11.6 (8.3) | 8.6 (6.7) | 0.14 |
| Polydrug use | 9.5 (6.5) | 10.7 (6.9) | 8.2 (5.8) | 0.14 |
| Injecting drug use, <i>n</i> (%) | 47 (84) | 24 (83) | 23 (85) | 0.55 |
| Injecting drug use, lifetime years: mean (s.d.) | 9.4 (7.7) | 10 (7.3) | 9 (8.2) | 0.65 |
| Overdose, lifetime number: mean (s.d.) | 5 (7) | 6 (8) | 4 (6.6) | 0.37 |
| Tried maintenance treatment, <i>n</i> (%) | 12 (21) | 7 (24) | 5 (18) | 0.75 |
| Detoxifications, lifetime number: mean (s.d.) | 5.8 (5.5) | 6 (6.1) | 5.7 (4.8) | 0.82 |
| Long-term residential stays, number: mean (s.d.) | 1.6 (1.5) | 1.8 (1.5) | 1.5 (1.6) | 0.56 |
| * <i>P</i> < 0.05. | | | | |