

Data supplement I Severity of suicidal ideation: univariate analyses

Variable	Passive ideation (n=185)		Active ideation (n=134)		Ideation with intent/plan (n=31)	
	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
Age group (years)						
18–29	1.18 (0.85–1.65)	0.32	0.87 (0.58–1.29)	0.48	1.58 (0.76–3.28)	0.22
31–44	0.93 (0.68–1.27)	0.64	1.34 (0.94–1.91)	0.11	1.06 (0.52–2.18)	0.87
45–59	1.05 (0.74–1.48)	0.80	0.94 (0.62–1.41)	0.76	0.69 (0.28–1.69)	0.41
> 59	0.49 (0.20–1.24)	0.12	0.41 (0.13–1.32)	0.12	0	
Ethnicity						
White	1.19 (0.85–1.65)	0.31	0.94 (0.64–1.40)	0.77	1.02 (0.46–2.23)	0.97
African American	1.53 (1.12–2.08)	0.007	1.42 (1.00–2.03)	0.05	1.01 (0.49–2.10)	0.97
Hispanic	0.50 (0.34–0.73)	0.0002	0.71 (0.47–1.06)	0.09	0.87 (0.40–1.91)	0.74
Male gender	1.20 (0.88–1.64)	0.24	1.29 (0.90–1.85)	0.17	1.87 (0.87–3.99)	0.10
New patient to ED	0.72 (0.47–1.10)	0.13	1.05 (0.67–1.64)	0.83	1.27 (0.54–2.99)	0.58
Shift during which survey was taken						
00.00–06.00	0.66 (0.43–1.03)	0.07	0.91 (0.57–1.46)	0.70	0.47 (0.14–1.56)	0.21
06.00–12.00	1.02 (0.76–1.65)	0.57	0.67 (0.23–1.94)	0.49	0.67 (0.23–1.94)	0.46
12.00–18.00	0.88 (0.62–1.25)	0.48	0.69 (0.46–1.06)	0.09	0.86 (0.38–1.94)	0.72
18.00–24.00	11.31 (0.96–1.79)	0.09	1.28 (0.89–1.84)	0.18	2.01 (0.99–4.10)	0.05
More than 1 ED visit in past 24 months	1.71 (1.24–2.35)	0.007	1.41 (0.98–2.01)	0.06	1.80 (0.86–3.79)	0.11
Any psychiatric ED visit within past 24 months	6.21 (3.58–10.77)	<0.0001	8.68 (4.94–15.25)	<0.0001	10.74 (4.58–25.21)	<0.0001
Presenting problem						
Cardiac	0.61 (0.22–1.72)	0.35	0.42 (0.10–1.74)	0.22	0	
Respiratory	1.02 (0.63–1.65)	0.94	1.17 (0.69–2.00)	0.56	0.85 (0.26–2.82)	0.79
Acute injury	0.96 (0.45–2.05)	0.93	1.20 (0.54–2.68)	0.65	0.71 (0.10–5.29)	0.73
Gastrointestinal/urinary	0.83 (0.56–1.21)	0.33	0.68 (0.43–1.08)	0.10	0.98 (0.42–2.30)	0.97
Pain	0.91 (0.55–1.50)	0.70	1.45 (0.88–2.40)	0.14	1.19 (0.41–3.44)	0.75
Infection	1.33 (0.87–2.03)	0.18	1.10 (0.66–1.84)	0.70	0.98 (0.34–2.83)	0.97
All other	1.26 (0.87–1.82)	0.23	1.05 (0.68–1.64)	0.82	1.45 (0.65–3.29)	0.36
Psychopathology						
Major depressive disorder	9.88 (7.02–13.80)	<0.0001	10.17 (6.85–15.09)	<0.0001	9.81 (4.35–22.12)	<0.0001
Bipolar disorder	9.59 (6.63–13.89)	<0.0001	9.97 (6.68–14.87)	<0.0001	12.63 (6.09–26.18)	<0.0001
Substance misuse	4.87 (3.45–6.88)	<0.0001	4.70 (3.19–6.91)	<0.0001	4.25 (2.03–8.89)	<0.0001
Anxiety	1.81 (0.99–3.30)	0.05	2.18 (1.14–4.15)	0.02	3.11 (1.06–9.13)	0.03
Panic	8.65 (6.11–12.25)	<0.0001	8.33 (5.68–12.23)	<0.0001	9.50 (4.60–19.60)	<0.0001
Traumatic stress symptoms	6.50 (4.43–9.54)	<0.0001	7.84 (5.20–11.84)	<0.0001	5.20 (2.40–11.28)	<0.0001
History of childhood sexual abuse	2.19 (1.56–3.07)	<0.0001	2.57 (1.7–3.77)	<0.0001	2.03 (0.95–4.35)	0.06

ED, emergency department.