

Supplementary Material to Burnett-order constitutive relations, second moment anisotropy and co-existing states in sheared dense gas-solid suspensions

Saikat Saha¹ and Meheboob Alam ^{1†}

¹Jawaharlal Nehru Centre for Advanced Scientific Research, Jakkur P.O., Bangalore 560064,
India

† Email address for correspondence: meheboob@jncasr.ac.in

Appendix A. Evaluation of collision integrals $\Theta_{\alpha\beta}$, $A_{\alpha\beta}$, $\hat{E}_{\alpha\beta}$ and $\hat{G}_{\alpha\beta}$

The collision integrals $\Theta_{\alpha\beta}$ (2.14), $A_{\alpha\beta}$ (2.16a), $\hat{E}_{\alpha\beta}$ (2.16b) and $\hat{G}_{\alpha\beta}$ (2.16c) appearing in the second-moment balance (2.24a-d) of the main text of Saha & Alam (2020) can be simplified to:

$$\left. \begin{aligned} 2\Theta_{xy} &= \frac{3(1+e)\rho\nu g_0 T}{\pi^{\frac{3}{2}}} \left(\cos 2\phi \mathcal{J}_{012}^{30} - \sin 2\phi \mathcal{J}_{102}^{30} \right) \\ \Theta_{xx} + \Theta_{yy} &= \frac{3(1+e)\rho\nu g_0 T}{\pi^{\frac{3}{2}}} \mathcal{J}_{002}^{30} \\ \Theta_{yy} - \Theta_{xx} &= \frac{3(1+e)\rho\nu g_0 T}{\pi^{\frac{3}{2}}} \left(\mathcal{J}_{102}^{30} \cos 2\phi + \mathcal{I}_{012}^{30} \sin 2\phi \right) \end{aligned} \right\}, \quad (\text{A } 1)$$

$$\left. \begin{aligned} A_{xx} &= -\frac{6(1-e^2)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left[\frac{1}{2} (\mathcal{H}_{003}^{30} - \cos 2\phi \mathcal{H}_{103}^{30} - \sin 2\phi \mathcal{H}_{013}^{30}) \right] \\ A_{yy} &= -\frac{6(1-e^2)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left[\frac{1}{2} (\mathcal{H}_{003}^{30} + \cos 2\phi \mathcal{H}_{103}^{30} + \sin 2\phi \mathcal{H}_{013}^{30}) \right] \\ A_{zz} &= -\frac{6(1-e^2)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \mathcal{H}_{003}^{12} \\ A_{xy} &= -\frac{6(1-e^2)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left[\frac{1}{2} (\cos 2\phi \mathcal{H}_{013}^{30} - \sin 2\phi \mathcal{H}_{103}^{30}) \right] \end{aligned} \right\}, \quad (\text{A } 2)$$

$$\left. \begin{aligned} \hat{E}_{xx} &= -(\hat{E}_{yy} + \hat{E}_{zz}), \\ \hat{E}_{yy} &= -\frac{6(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \\ &\quad \times \left[\eta \left\{ \mathcal{H}_{101}^{31} - \mathcal{H}_{011}^{32} + \cos 2\phi \left(\mathcal{H}_{201}^{31} - \mathcal{H}_{021}^{31} + \mathcal{H}_{111}^{50} \right) - \sin 2\phi \left(\mathcal{H}_{201}^{30} + \mathcal{H}_{021}^{32} \right) \right\} \right. \\ &\quad \left. + 3\lambda^2 \left\{ \mathcal{H}_{001}^{32} + \cos 2\phi \left(\mathcal{H}_{101}^{32} + \mathcal{H}_{011}^{31} \right) + \sin 2\phi \left(\mathcal{H}_{011}^{32} - \mathcal{H}_{101}^{31} \right) \right\} \right] \\ \hat{E}_{zz} &= -\frac{12(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left\{ -\eta \left(\mathcal{H}_{101}^{31} - \mathcal{H}_{011}^{32} \right) - \lambda^2 \left(\mathcal{H}_{001}^{32} \right) \right\} \\ \hat{E}_{xy} &= -\frac{12(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left[\frac{1}{2} \eta \left\{ \left(2\mathcal{H}_{111}^{31} - \mathcal{H}_{021}^{32} - \mathcal{H}_{201}^{30} \right) \cos 2\phi \right. \right. \\ &\quad \left. \left. + \left(\mathcal{H}_{021}^{31} + \mathcal{H}_{111}^{32} - \mathcal{H}_{111}^{30} - \mathcal{H}_{201}^{31} \right) \sin 2\phi \right\} \right. \\ &\quad \left. + \frac{3}{2} \lambda^2 \left\{ \left(\mathcal{H}_{011}^{32} - \mathcal{H}_{101}^{31} \right) \cos 2\phi - \left(\mathcal{H}_{101}^{32} + \mathcal{H}_{011}^{31} \right) \sin 2\phi \right\} \right] \end{aligned} \right\} \quad (\text{A } 3)$$

$$\left. \begin{aligned}
\hat{G}_{xx} &= -(\hat{G}_{yy} + \hat{G}_{zz}) \\
\hat{G}_{yy} &= -\frac{6(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} 2R \left\{ \mathcal{K}_{00}^{31} + \cos 2\phi \left(\mathcal{K}_{10}^{31} + \mathcal{K}_{01}^{30} \right) + \sin 2\phi \left(\mathcal{K}_{01}^{31} - \mathcal{K}_{10}^{30} \right) \right\} \\
\hat{G}_{zz} &= -\frac{6(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left\{ -4R\mathcal{K}_{00}^{31} \right\} \\
\hat{G}_{xy} &= -\frac{6(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} (-4R) \left\{ \frac{1}{2} \cos 2\phi \left(\mathcal{K}_{01}^{31} - \mathcal{K}_{10}^{30} \right) + \frac{1}{2} \sin 2\phi \left(\mathcal{K}_{10}^{31} + \mathcal{K}_{01}^{30} \right) \right\}
\end{aligned} \right\} \quad (\text{A } 4)$$

In the above expressions, $\mathcal{H}_{\alpha\beta\gamma}^{\delta p}$, $\mathcal{J}_{\alpha\beta\gamma}^{\delta p}$ and $\mathcal{K}_{\alpha\beta}^{\delta p}$ represent tensorial integrals defined as

$$\begin{aligned}
\mathcal{H}_{\alpha\beta\gamma}^{\delta p}(\eta, R, \phi, \lambda) &\equiv \int_{\theta=0}^{2\pi} \int_{\varphi=0}^{\pi} \sin^{\alpha} 2\theta \cos^{\beta} 2\theta \sin^{\delta} \varphi \cos^p \varphi \\
&\quad \times (1 - \eta \sin^2 \varphi \cos 2\theta + \lambda^2 (3 \sin^2 \varphi - 2))^{\frac{\gamma}{2}} \mathfrak{F}(\chi[\eta, R, \phi, \lambda; \theta, \varphi]) d\varphi d\theta, \quad (\text{A } 5)
\end{aligned}$$

$$\begin{aligned}
\mathcal{J}_{\alpha\beta\gamma}^{\delta p}(\eta, R, \phi, \lambda) &\equiv \int_{\theta=0}^{2\pi} \int_{\varphi=0}^{\pi} \sin^{\alpha} 2\theta \cos^{\beta} 2\theta \sin^{\delta} \varphi \cos^p \varphi \\
&\quad \times \{1 - \eta \sin^2 \varphi \cos 2\theta + \lambda^2 (3 \sin^2 \varphi - 2)\}^{\frac{\gamma}{2}} \mathfrak{G}(\chi[\eta, R, \phi, \lambda; \theta, \varphi]) d\varphi d\theta, \quad (\text{A } 6)
\end{aligned}$$

$$\begin{aligned}
\mathcal{K}_{\alpha\beta}^{\delta p}(\eta, R, \phi, \lambda) &\equiv \int_{\theta=0}^{2\pi} \int_{\varphi=0}^{\pi} \sin^{\alpha} 2\theta \cos^{\beta} 2\theta \sin^{\delta} \varphi \cos^p \varphi [(1 - 2\lambda^2) \{\sin(2\phi + 2\theta) - \cos \varphi \\
&\quad \times \cos(2\phi + 2\theta)\} + \sin^2 \varphi \{3\lambda^2 \sin(2\phi + 2\theta) - \eta \sin 2\phi\}] \mathfrak{G}(\chi[\eta, R, \phi, \lambda; \theta, \varphi]) d\varphi d\theta, \quad (\text{A } 7)
\end{aligned}$$

where $\mathfrak{F}(\chi)$ and $\mathfrak{G}(\chi)$ are given by (2.18a) and (2.18b), respectively, and each integral is evaluated over $\theta \in (0, 2\pi)$ and $\varphi \in (0, \pi)$; see figure 1 for the definitions of θ and φ with reference to the contact vector \mathbf{k} . We follow an analytical approach to evaluate the above elliptic integrals following our previous works (Saha & Alam 2014, 2016) on sheared dry granular flows – the underlying methodology is identical to above works and is briefly discussed below.

Since both $\mathfrak{F}(\chi)$ and $\mathfrak{G}(\chi)$ are analytical functions of χ , we substitute (2.19) into (2.18a) and (2.18b) and use the power-series representations for the complementary error function and the exponential. This yields the following infinite series representation (Saha & Alam 2014) for $\mathfrak{F}(\chi)$ and $\mathfrak{G}(\chi)$:

$$\begin{aligned}
\mathfrak{F}(\eta, R, \phi, \lambda; \theta, \varphi) &= -\sqrt{\pi} \left[\frac{3R \sin^2 \varphi \cos(2\phi + 2\theta)}{\{1 - \eta \sin^2 \varphi \cos 2\theta + \lambda^2 (3 \sin^2 \varphi - 2)\}^{\frac{1}{2}}} \right. \\
&\quad \left. + \left\{ \frac{2R \sin^2 \varphi \cos(2\phi + 2\theta)}{\{1 - \eta \sin^2 \varphi \cos 2\theta + \lambda^2 (3 \sin^2 \varphi - 2)\}^{\frac{1}{2}}} \right\}^3 \right] \\
&\quad + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \frac{3}{(2n-1)(2n-3)} \left[\frac{2R \sin^2 \varphi \cos(2\phi + 2\theta)}{\{1 - \eta \sin^2 \varphi \cos 2\theta + \lambda^2 (3 \sin^2 \varphi - 2)\}^{\frac{1}{2}}} \right]^{2n}, \quad (\text{A } 8)
\end{aligned}$$

$$\begin{aligned}
\mathfrak{G}(\eta, R, \phi, \lambda; \theta, \varphi) &= \sqrt{\pi} \left[\frac{1}{2} + \frac{4R^2 \sin^4 \varphi \cos^2(2\phi + 2\theta)}{1 - \eta \sin^2 \varphi \cos 2\theta + \lambda^2 (3 \sin^2 \varphi - 2)} \right] \\
&\quad + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \frac{2}{(2n-1)(2n+1)} \left[\frac{2R \sin^2 \varphi \cos(2\phi + 2\theta)}{\{1 - \eta \sin^2 \varphi \cos 2\theta + \lambda^2 (3 \sin^2 \varphi - 2)\}^{\frac{1}{2}}} \right]^{2n+1} \quad (\text{A } 9)
\end{aligned}$$

Substituting (A 8-A 9) into (A 5-A 7) and carrying out term-by-term integrations over




FIGURE 1. Sketch of the coordinate system (x, y, z) , with the shear-field $[u_x(y) = \dot{\gamma}y]$ being applied on the (x, y) -plane and z is the mean vorticity direction; adapted from Saha & Alam (2016). The eigen-directions of the strain-rate tensor \mathbf{D} are marked as $(|D_1\rangle, |D_2\rangle, |D_3\rangle)$ and that of the second moment tensor \mathbf{M} as $(|M_1\rangle, |M_2\rangle, |M_3\rangle)$. Note that \mathbf{k} represents the contact vector between two colliding particles, that makes an angle φ with z -direction and its projection on the (x, y) shear-plane makes an angle θ with $|M_1\rangle$ of \mathbf{M} . The *principle axes frame* (x', y', z') is given by the orthonormal triad of $(|M_1\rangle, |M_2\rangle, |M_3\rangle)$ which represents a rotation of the original (x, y, z) axes by an angle $\vartheta = \phi + \pi/4$ about the z -axis.

$\theta \in (0, 2\pi)$ and $\varphi \in (0, \pi)$, we obtain infinite series expressions for $\mathcal{H}_{\alpha\beta\gamma}^{\delta p}$, $\mathcal{J}_{\alpha\beta\gamma}^{\delta p}$ and $\mathcal{K}_{\alpha\beta\gamma}^{\delta p}$ as functions of η , $\sin \phi$, λ and R .

Truncating the infinite series representations of $\mathcal{H}_{\alpha\beta\gamma}^{\delta p}$, $\mathcal{J}_{\alpha\beta\gamma}^{\delta p}$ and $\mathcal{K}_{\alpha\beta\gamma}^{\delta p}$ at the Burnett order [i.e. the second-order in $O(\eta^i \lambda^j R^k \sin^l 2\phi)$, with $i + j + k + l \leq 2$], the following Burnett-order expressions are obtained:

$$\frac{1}{2} \left(\cos 2\phi \mathcal{J}_{012}^{30} - \sin 2\phi \mathcal{J}_{102}^{30} \right) = -\frac{4\pi}{15} \left(8R + \sqrt{\pi}\eta \cos 2\phi \right), \quad (\text{A } 10)$$

$$\mathcal{J}_{002}^{30} = \frac{4\pi}{105} \left\{ \sqrt{\pi} \left(35 + 96R^2 + 14\lambda^2 \right) + 48R\eta \cos 2\phi \right\} \quad (\text{A } 11)$$

$$(\mathcal{J}_{102}^{30} \cos 2\phi + \mathcal{I}_{012}^{30} \sin 2\phi) = -\frac{8}{15} \pi^{\frac{3}{2}} \eta \sin 2\phi, \quad (\text{A } 12)$$

$$\begin{aligned} & \left[\frac{1}{2} \left(\mathcal{H}_{003}^{30} - \cos 2\phi \mathcal{H}_{103}^{30} - \sin 2\phi \mathcal{H}_{013}^{30} \right) \right] \\ &= \frac{2\pi}{105} \left(70 + 9\eta^2 + 42\lambda^2 + 288R^2 + 72\sqrt{\pi}\eta R \cos 2\phi + 42\eta \sin 2\phi \right), \quad (\text{A } 13) \end{aligned}$$

$$\begin{aligned} & \left[\frac{1}{2} \left(\mathcal{H}_{003}^{30} + \cos 2\phi \mathcal{H}_{103}^{30} + \sin 2\phi \mathcal{H}_{013}^{30} \right) \right] \\ &= \frac{2\pi}{105} \left(70 + 9\eta^2 + 42\lambda^2 + 288R^2 + 72\sqrt{\pi}\eta R \cos 2\phi - 42\eta \sin 2\phi \right), \quad (\text{A } 14) \end{aligned}$$

$$\mathcal{H}_{003}^{12} = \frac{2\pi}{105} \left(70 + 3\eta^2 - 84\lambda^2 + 96R^2 + 24\sqrt{\pi}\eta R \cos 2\phi \right), \quad (\text{A } 15)$$

$$\left[\frac{1}{2} \left(\cos 2\phi \mathcal{H}_{013}^{30} - \sin 2\phi \mathcal{H}_{103}^{30} \right) \right] = -\frac{4\pi}{5} \left(2\sqrt{\pi}R + \eta \cos 2\phi \right), \quad (\text{A } 16)$$

$$\widehat{E}_{yy} = -\frac{6(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left[\frac{8\pi}{105} \left(\eta^2 + 21\lambda^2 + 6\sqrt{\pi}\eta R \cos 2\phi - 21\eta \sin 2\phi \right) \right], \quad (\text{A } 17)$$

$$\widehat{E}_{zz} = -\frac{12(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left[-\frac{8\pi}{105} \left(\eta^2 + 21\lambda^2 + 6\sqrt{\pi}\eta R \cos 2\phi \right) \right], \quad (\text{A } 18)$$

$$\widehat{E}_{xy} = -\frac{12(1+e)\rho\nu g_0 T^{\frac{3}{2}}}{\sigma\pi^{\frac{3}{2}}} \left[-\frac{4\pi}{5} \eta \cos 2\phi \right], \quad (\text{A } 19)$$

$$2R \left\{ \mathcal{K}_{00}^{31} + \cos 2\phi \left(\mathcal{K}_{10}^{31} + \mathcal{K}_{01}^{30} \right) + \sin 2\phi \left(\mathcal{K}_{01}^{31} - \mathcal{K}_{10}^{30} \right) \right\} = \frac{128\pi R^2}{105}, \quad \mathcal{K}_{00}^{31} = \frac{64\pi R}{105}, \quad (\text{A } 20)$$

$$\frac{1}{2} \cos 2\phi \left(\mathcal{K}_{01}^{31} - \mathcal{K}_{10}^{30} \right) + \frac{1}{2} \sin 2\phi \left(\mathcal{K}_{10}^{31} + \mathcal{K}_{01}^{30} \right) = \frac{2\pi}{105} \left[\sqrt{\pi} \left(21 + 32R^2 \right) - 16\eta R \cos 2\phi \right]. \quad (\text{A } 21)$$

A.1. Stress tensor up-to fourth-order

The diagonal components of the dimensionless stress tensor, $P_{ii}^* = \tilde{P}_{ii}/\rho_p\nu U_R^2$, with $U_R = \dot{\gamma}\sigma/2$, correct up-to $\mathcal{O}(\eta^i\lambda^j R^k \sin^l 2\phi, i+j+k+l \leq 4)$, have following expressions:

$$\begin{aligned} \frac{P_{xx}^*}{T^*} &= (1 + \lambda^2 + \eta \sin 2\phi) + \frac{2(1+e)\nu g_0}{35} \left[(35 + 96R^2 + 14\eta \sin 2\phi + 14\lambda^2) \right. \\ &\quad \left. + \frac{8}{\sqrt{\pi}} \eta R \cos 2\phi \left\{ 3(66 + 5\eta^2 - 22\lambda^2) - 160R^2 - 22\eta \sin 2\phi \right\} \right], \end{aligned} \quad (\text{A } 22)$$

$$\begin{aligned} \frac{P_{yy}^*}{T^*} &= (1 + \lambda^2 - \eta \sin 2\phi) + \frac{2(1+e)\nu g_0}{35} \left[(35 + 96R^2 - 14\eta \sin 2\phi + 14\lambda^2) \right. \\ &\quad \left. + \frac{8}{\sqrt{\pi}} \eta R \cos 2\phi \left\{ 3(66 + 5\eta^2 - 22\lambda^2) - 160R^2 + 22\eta \sin 2\phi \right\} \right], \end{aligned} \quad (\text{A } 23)$$

$$\begin{aligned} \frac{P_{zz}^*}{T^*} &= (1 - 2\lambda^2) + \frac{2(1+e)\nu g_0}{35} \left[(35 + 32R^2 - 28\lambda^2) \right. \\ &\quad \left. + \frac{8}{\sqrt{\pi}} \eta R \cos 2\phi (66 + 3\eta^2 - 32R^2) \right]. \end{aligned} \quad (\text{A } 24)$$

where $T^* = \tilde{T}/U_R^2$ is the dimensionless temperature.

To calculate the fourth-order viscosity, we need to retain terms up-to the fifth-order $\mathcal{O}(\eta^i\lambda^j R^k \sin^l 2\phi, i+j+k+l \leq 5)$ in the shear stress. The dimensionless shear stress, $P_{xy}^* = \tilde{P}_{xy}/\rho_p\nu U_R^2$, with $U_R = \dot{\gamma}\sigma/2$, at $\mathcal{O}(R^5)$, can be written as :

$$\begin{aligned} \frac{P_{xy}^*}{T^*} &= -\eta \cos 2\phi - \frac{4(1+e)\nu g_0}{105\sqrt{\pi}} \left[21R \left\{ 8 + \sqrt{\pi} \frac{\eta \cos 2\phi}{R} \right\} \right. \\ &\quad \left. + 4R^3 \left\{ 32 + 12 \frac{\lambda^2}{R^2} - \frac{\eta^2}{R^2} (2 + \cos 4\phi) \right\} \right. \\ &\quad \left. + \frac{R^5}{143} \left\{ -5120 + 15 \frac{\eta^2}{R^2} \left(64 - 5 \frac{\eta^2}{R^2} \right) (3 + 2 \cos 4\phi) \right. \right. \\ &\quad \left. \left. + 52 \frac{\lambda^2}{R^2} \left(-128 - 33 \frac{\lambda^2}{R^2} + 12 (2 + \cos 4\phi) \frac{\eta^2}{R^2} \right) \right\} \right]. \end{aligned} \quad (\text{A } 25)$$

Appendix B. Expressions for coefficients a_i in Eqn. (3.1)

The expressions for the coefficients of the temperature equation (3.1) are given by:

$$a_0 = 10000(1+e)^5\pi^{3/2}St_d^3\{63(1-e)\pi - 4(11+e)St_d\}^2(4+St_d^2)\nu^3g_0^3 \quad (B\ 1)$$

$$\begin{aligned} a_1 = & -2000(1+e)^2\pi St_d\nu g_0 \left(18522000(1-e)\pi^2 + 463050(1-e)\pi^3 St_d \{5 - 2(1+e)(1-3e)\nu g\} \right. \\ & + 29400\pi^2 St_d^2 \{5(67-55e) + 2(1+e)(23+4e-3e^2)\nu g\} \\ & + 441\pi^2 St_d^3 [125(13-9e)\pi - 50(1+e)(11-54e+27e^2)\pi\nu g \\ & + 40(1+e)^2(1+3e)\{6(1-e) - \pi(1-3e)\}\nu^2g_0^2 - 864(1-e)^2(3-e)(1+e)^4\nu^3g_0^3] \\ & + 56\pi St_d^4 [1750(13-9e)\pi + 25(1+e)(277+54e-27e^2)\pi\nu g \\ & - 20(1+e)^2(1+3e)(6(11+e) - (17+3e)\pi)\nu^2g_0^2 + 648(1-e)(3-e)(1+e)^4(11+e)\nu^3g_0^3] \\ & + 12(1+e)^2 St_d^5 \nu^2 g_0^2 [2205(1-e)(1+3e)\pi^2 \\ & - (3-e)(1+e)^2 \{64(11+e)^2 + 3969(1-e)^2\pi^2\}\nu g] \\ & \left. - 336(1+e)^2(11+e)\pi St_d^6 \nu^2 g_0^2 \{5(1+3e) - 9(1-e)(3-e)(1+e)^2\nu g\} \right) \quad (B\ 2) \end{aligned}$$

$$\begin{aligned} a_2 = & -600(1+e)\sqrt{\pi} St_d\nu g_0 \left(54022500(1-e)(1+3e)\pi^3 \right. \\ & + 444528000(1-e)(3-e)(1+e)^2\pi^2 St_d\nu g + 14700\pi^3 St_d^2 \{175(11-9e)(1+3e) \\ & + 5(1+e)(1829-381e-1773e^2+549e^3)\nu g + 126(1-3e)(1-e)(1+e)^3(-29+9e)\nu^2g_0^2\} \\ & + 8400(1+e)^2\pi^2 St_d^3 \nu g_0 \{5(11399-13078e+3071e^2) \\ & + 14(1+e)(541-45e-133e^2+21e^3)\nu g\} + 147\pi^2 St_d^4 [1875(13-9e)(1+3e)\pi \\ & + 125(1+e)(1351+237e-999e^2+243e^3)\pi\nu g - 50(1+e)^2\{16(1+3e)^2 \\ & + (1117-4836e-1566e^2+3636e^3-783e^4)\pi\}\nu^2g_0^2 \\ & + 1440(3-e)(1+e)^4(1+3e)\{6(1-e) - \pi(1-3e)\}\nu^3g_0^3 \\ & - 15552(1-e)^2(3-e)^2(1+e)^6\nu^4g_0^4] + 28(1+e)^2\pi St_d^5 \nu g_0 [125(705-221e)(13-9e)\pi \\ & + 50(1+e)(6417-17e-1689e^2+153e^3)\pi\nu g \\ & - 320(3-e)(1+e)^2(1+3e)\{6(11+e) - (17+3e)\pi\}\nu^2g_0^2 \\ & + 5184(1-e)(3-e)^2(1+e)^4(11+e)\nu^3g_0^3] - 24(1+e)^2 St_d^6 \nu^2 g_0^2 [1225(1+3e)^2\pi^2 \\ & - 4410(3-e)(1-e)(1+e)^2(1+3e)\pi^2\nu g_0 + (3-e)^2(1+e)^4\{64(11+e)^2 \\ & + 3969(1-e)^2\pi^2\}\nu^2g_0^2] \left. \right) \quad (B\ 3) \end{aligned}$$

$$\begin{aligned} a_3 = & +2520\pi \left(96468750(1-e)\pi^3 + 91875\pi^2 St_d^2 [25(13-9e)\pi + 80(1+e)(3-8e+6e^2)\pi\nu g_0 \right. \\ & - 28(1-e)(1+e)^2\{36(3-e)(1+3e) + (1-3e)^2\pi\}\nu^2g_0^2] \\ & - 317520000(1-e)(3-e)^2(1+e)^4\pi St_d^3 \nu^3 g_0^3 \\ & + 350(1+e)\pi^2 St_d^4 \nu g_0 [125(11-24e)(13-9e)\pi \\ & - 25(1+e)\{12(1+3e)(937-1073e+252e^2) + (47-645e+1485e^2-567e^3)\pi\}\nu g_0 \\ & - 20(1+e)^2\{6(5333-2151e-4815e^2+2991e^3-462e^4) + 7(1-3e)^2(1+3e)\pi\}\nu^2g_0^2 \\ & + 12096(1-3e)(7-2e)(1-e)(3-e)(1+e)^4\nu^3g_0^3] \\ & - 800(3-e)(1+e)^4\pi St_d^5 \nu^3 g_0^3 [5(51851-59290e+13775e^2) \\ & + 126(1+e)(5+3e)(53-32e+3e^2)\nu g_0] \\ & - 42(1+e)^2\pi St_d^6 \nu^2 g_0^2 [625(13-9e)(19-6e)(1+3e)\pi \\ & + 250(1+e)(1976+129e-1557e^2+615e^3-75e^4)\pi\nu g_0 \\ & \left. - 100(3-e)(1+e)^2\{16(1+3e)^2 + (523-2316e-882e^2+1692e^3-297e^4)\pi\}\nu^2g_0^2 \right) \end{aligned}$$

$$\begin{aligned}
& +1440(3-e)^2(1+e)^4(1+3e)\{6(1-e)-(1-3e)\pi\}\nu^3g_0^3 \\
& -10368(1-e)^2(3-e)^3(1+e)^6\nu^4g_0^4] \\
& -8(3-e)(1+e)^4St_d^7\nu^3g_0^3[125(4709-4790e+977e^2)\pi \\
& +50(1+e)(3093+443e-1149e^2+45e^3)\pi\nu g_0 \\
& -160(3-e)(1+e)^2(1+3e)\{6(11+e)-(17+3e)\pi\}\nu^2g_0^2 \\
& +1728(1-e)(3-e)^2(1+e)^4(11+e)\nu^3g_0^3]) \quad (\text{B } 4)
\end{aligned}$$

$$\begin{aligned}
a_4 = & +63(1+e)\pi^{3/2}St_d\nu g_0 \left(1543500000(59-23e)(1-e)\pi^2 \right. \\
& +52500\pi St_d^2[25(14647-16830e+4919e^2)\pi \\
& +140(1+e)(2041-6273e+5999e^2-1383e^3)\pi\nu g_0 \\
& -588(1-e)(1+e)^2\{360(3-e)^2(1+3e)+(1-3e)^2(59-19e)\pi\}\nu^2g_0^2] \\
& -20321280000(1-e)(3-e)^3(1+e)^4St_d^3\nu^3g_0^3-175\pi St_d^4[625(13-9e)(107+193e)\pi \\
& -500(1+e)(28181-94445e+71523e^2-14283e^3)\pi\nu g_0 \\
& +100(1+e)^2\{96(3-e)(1+3e)(4267-4873e+1134e^2) \\
& +(10223-137768e+356922e^2-219384e^3+36855e^4)\pi\}\nu^2g_0^2 \\
& +5760(3-e)(1+e)^3\{6(1736-439e-1557e^2+831e^3-123e^4) \\
& +7(1-3e)^2(1+3e)\pi\}\nu^3g_0^3-290304(27-7e)(1-3e)(1-e)(3-e)^2(1+e)^5\nu^4g_0^4] \\
& -38400(3-e)^2(1+e)^4St_d^5\nu^3g_0^3\{5(34879-39854e+9151e^2) \\
& +42(1+e)(519+41e-191e^2+15e^3)\nu g_0\} \\
& -672(3-e)(1+e)^2St_d^6\nu^2g_0^2[625(1+3e)(382-391e+81e^2)\pi \\
& +375(1+e)(1303+394e-984e^2+198e^3-15e^4)\pi\nu g_0 \\
& -150(3-e)(1+e)^2\{16(1+3e)^2+(325-1476e-654e^2+1044e^3-135e^4)\pi\}\nu^2g_0^2 \\
& +1440(3-e)^2(1+e)^4(1+3e)\{6(1-e)-(1-3e)\pi\}\nu^3g_0^3 \\
& \left. -7776(1-e)^2(3-e)^3(1+e)^6\nu^4g_0^4\right] \quad (\text{B } 5)
\end{aligned}$$

$$\begin{aligned}
a_5 = & +3780(3-e)(1+e)^2\pi St_d^2\nu^2g_0^2 \left(463050000(11-5e)(1-e)\pi^2 \right. \\
& +4200\pi St_d^2[25(6405-8420e+3047e^2)\pi \\
& +140(1+e)(963-3040e+2881e^2-660e^3)\pi\nu g_0 \\
& -588(1-e)(1+e)^2\{120(3-e)^2(1+3e)+(29-9e)(1-3e)^2\pi\}\nu^2g_0^2] \\
& -304819200(1-e)(3-e)^3(1+e)^4St_d^3\nu^3g_0^3-7\pi St_d^4[625(13-9e)(131+205e)\pi \\
& -500(1+e)(12821-45203e+33483e^2-6381e^3)\pi\nu g_0 \\
& +100(1+e)^2\{48(3-e)(1+3e)(2873-3281e+756e^2) \\
& +(5147-66416e+173322e^2-104688e^3+16443e^4)\pi\}\nu^2g_0^2 \\
& +2880(3-e)(1+e)^3\{2(2556-235e-2253e^2+963e^3-135e^4) \\
& +7(1-3e)^2(1+3e)\pi\}\nu^3g_0^3-145152(1-3e)(13-3e)(1-e)(3-e)^2(1+e)^5\nu^4g_0^4] \\
& -768(3-e)^2(1+e)^4St_d^5\nu^3g_0^3\{5(8783-10054e+2279e^2) \\
& \left. +42(1+e)(127+21e-55e^2+3e^3)\nu g_0\} \right) \quad (\text{B } 6)
\end{aligned}$$

$$\begin{aligned}
a_6 = & +15876(3-e)^2(1+e)^3\sqrt{\pi}St_d^3\nu^3g_0^3\Big(132300000(17-9e)(1-e)\pi^2 \\
& +St_d^2[5000(32639-50458e+21923e^2)\pi^2 \\
& +252000(1+e)(605-1967e+1843e^2-417e^3)\pi^2\nu g_0 \\
& -352800(1-e)(1+e)^2\pi\{180(3-e)^2(1+3e)+(57-17e)(1-3e)^2\pi\}\nu^2g_0^2] \\
& -34836480(1-e)(3-e)^3(1+e)^4St_d^3\nu^3g_0^3 \\
& -\pi St_d^4[625(1951+1362e-1969e^2)\pi \\
& -500(1+e)(7757-28985e+20859e^2-3663e^3)\pi\nu g_0 \\
& +100(1+e)^2\{96(3-e)(1+3e)(724-829e+189e^2) \\
& +(3455-42632e+112122e^2-66456e^3+9639e^4)\pi\}\nu^2g_0^2 \\
& +1920(3-e)(1+e)^3\{3(1009+78e-864e^2+258e^3-33e^4) \\
& +7(1-3e)^2(1+3e)\pi\}\nu^3g_0^3-145152(1-3e)(1-e)(3-e)^2(5-e)(1+e)^5\nu^4g_0^4]\Big) \quad (B7)
\end{aligned}$$

$$\begin{aligned}
a_7 = & 3048192(3-e)^3(1+e)^4St_d^4\nu^4g_0^4\Big(275625(47-29e)(1-e)\pi^2 \\
& +5\pi St_d^2[25(3325-6284e+3211e^2)\pi+210(1+e)(427-1434e+1325e^2-294e^3)\pi\nu g_0 \\
& -1764(1-e)(1+e)^2\{18(3-e)^2(1+3e)+(1-3e)^2(7-2e)\pi\}\nu^2g_0^2] \\
& -36288(1-e)(3-e)^3(1+e)^4St_d^3\nu^3g_0^3\Big) \quad (B8)
\end{aligned}$$

$$\begin{aligned}
a_8 = & +2286144(1-e)(3-e)^4(1+e)^5\sqrt{\pi}St_d^5\nu^5g_0^5\Big(264600(43-31e)\pi \\
& +St_d^2[25(4069-5917e)\pi+420(1+e)(321-796e+219e^2)\pi\nu g_0 \\
& -1764(1+e)^2\{24(3-e)^2(1+3e)+(1-3e)^2(11-3e)\pi\}\nu^2g_0^2]\Big) \quad (B9)
\end{aligned}$$

$$a_9 = +725896442880(13-11e)(1-e)(3-e)^5(1+e)^6\pi St_d^6\nu^6g_0^6 \quad (B10)$$

$$a_{10} = +1451792885760(3-e)^6(1-e)^2(1+e)^7\sqrt{\pi}St_d^7\nu^7g_0^7. \quad (B11)$$

Note that $a_{10} = 0 = a_9 = a_8$ for $e = 1$, leading to a seventh-order polynomial for the temperature equation (3.1).

B.1. Coefficients in equation (3.2)

The expressions for the coefficients of the quadratic polynomial (3.2) are given by

$$\begin{aligned}
\mathbf{a} = & \Big\{4375(403507877743-2246328611260e+3804527790618e^2 \\
& -2609043261628e^3+637175564527e^4)+7000(1841694494389-7655913090100e \\
& +9936315870558e^2-5002578505972e^3+804447129877e^4)\nu g_0 \\
& -1600(62781766140229-251533031265556e+383366453401950e^2 \\
& -256170205851796e^3+62747567827717e^4)\nu^2g_0^2-5120(15714866428013 \\
& -15498531439412e-53893773596274e^2+89513065956556e^3-35535888468883e^4)\nu^3g_0^3 \\
& +16384(55881178513091-252823502621588e+422083141225338e^2 \\
& -306989989576772e^3+81830578488731e^4)\nu^4g_0^4\Big\}, \quad (B12)
\end{aligned}$$

$$\begin{aligned}
\mathbf{b} = & -\left\{ 625(18778586124871 - 207069149476948e + 405023838452178e^2 \right. \\
& - 286513612682692e^3 + 68321253421567e^4) - 4000(195746263524613 \\
& - 832915077870466e + 1329037811913048e^2 - 934639544844094e^3 \\
& + 243370589968483e^4)\nu g_0 - 3200(1463725651004753 - 5379389667615500e \\
& + 7142510907700158e^2 - 4045727120864924e^3 + 816845472782489e^4)\nu^2 g_0^2 \\
& + 10240(1346413690308923 - 5798356073502980e + 9330745323357066e^2 \\
& - 6630820234286324e^3 + 1752390117615923e^4)\nu^3 g_0^3 \\
& + 262144(16659939237511 - 30369558221428e - 11077909546134e^2 \\
& \left. + 44890628459564e^3 - 20186596004777e^4)\nu^4 g_0^4 \right\}, \tag{B 13}
\end{aligned}$$

$$\begin{aligned}
\mathbf{c} = & 48\left\{ 1875(5292277916157 - 7198799909266e - 8983084338480e^2 \right. \\
& + 18332139150594e^3 - 7387862080253e^4) - 1000(653424923400545 \\
& - 2603969210032424e + 3825404450727618e^2 - 2459579042010032e^3 \\
& + 584400338366645e^4)\nu g_0 + 2400(449794922964867 - 2000985299639524e \\
& + 3319509658053858e^2 - 2435905168315812e^3 + 667705483931683e^4)\nu^2 g_0^2 \\
& + 7680(1001122119985027 - 3916700322857816e + 5710226196969174e^2 \\
& - 3680145930163120e^3 + 885453528512687e^4)\nu^3 g_0^3 \\
& + 4096(1 - e)(1554567774124823 - 5288085618364101e \\
& \left. + 5986900650648069e^2 - 2247667963924055e^3)\nu^4 g_0^4 \right\}. \tag{B 14}
\end{aligned}$$

Appendix C. Ignited state: analysis in the principal axes frame

The second-moment balance equation (4.1) in the principal-axes frame can be written as four independent equations (Alam, Saha & Gupta 2019):

$$\begin{aligned}
-4\eta\rho T\dot{\gamma} \cos 2\phi + 2\dot{\gamma}[(\Theta_{x'x'} - \Theta_{y'y'}) \cos 2\phi - 2\Theta_{x'y'} \sin 2\phi] + \frac{6\rho T\dot{\gamma}}{St_d} &= A_{\alpha'\alpha'}, \tag{C 1} \\
-4\eta\rho T\dot{\gamma} \cos 2\phi + 2\dot{\gamma}[(\Theta_{x'x'} - \Theta_{y'y'}) \cos 2\phi - 2\Theta_{x'y'} \sin 2\phi] + \frac{12\rho T\dot{\gamma}\lambda^2}{St_d} &= -3\hat{\Gamma}_{z'z'}, \tag{C 2} \\
4(1 + \lambda^2)\rho T\dot{\gamma} \cos 2\phi + 2\dot{\gamma}(\Theta_{x'x'} + \Theta_{y'y'}) \cos 2\phi - \frac{4\rho T\dot{\gamma}\eta}{St_d} &= \Gamma_{x'x'} - \Gamma_{y'y'}, \tag{C 3} \\
2\rho T\dot{\gamma}[\eta - (1 + \lambda^2) \sin 2\phi] - (\Theta_{x'x'} + \Theta_{y'y'})\dot{\gamma} \sin 2\phi &= \Gamma_{x'y'}. \tag{C 4}
\end{aligned}$$

They represent (i) the trace of (4.1), with $A_{\alpha'\alpha'} = A_{x'x'} + A_{y'y'} + A_{z'z'}$, (ii) the z' - z' component of the deviatoric part of (4.1) (iii) the difference between the x' - x' and y' - y' components and (iv) the off-diagonal x' - y' component of (4.1). The expressions for various integrals $A_{\alpha'\beta'}$, $\hat{E}_{\alpha'\beta'}$, $\hat{G}_{\alpha'\beta'}$ and $\Theta_{\alpha'\beta'}$ appearing in (C 1-C 4) remain the same as those for the dry granular flow; these have been evaluated explicitly by Saha & Alam (2016).

Retaining terms up-to fourth-order $O(\eta^i \lambda^j R^k \sin^l 2\phi)$, $i+j+k+l \leq 4$ in the expressions of $A_{\alpha'\beta'}$, $\hat{E}_{\alpha'\beta'}$, $\hat{G}_{\alpha'\beta'}$ and $\Theta_{\alpha'\beta'}$, the above equations (C 1-C 4) simplify to the following

set of four coupled nonlinear algebraic equations

$$\begin{aligned} & 5\sqrt{\pi}\eta R \cos 2\phi + 4(1+e)\nu g_0 R \left(\sqrt{\pi}\eta \cos 2\phi + 8R \right) \\ & - \frac{3}{4}(1-e^2)\nu g_0 (10 + \eta^2 + 32R^2 + 8\sqrt{\pi}\eta R \cos 2\phi) - \frac{15\sqrt{\pi}R}{St_d} \\ & - \frac{1}{28}\nu g_0 R^4 \left\{ 256 - 16(2 + \cos 4\phi)\left(\frac{\eta}{R}\right)^2 + 192\left(\frac{\lambda}{R}\right)^2 - 6\left(\frac{\eta}{R}\right)^2\left(\frac{\lambda}{R}\right)^2 + \frac{3}{4}\left(\frac{\eta}{R}\right)^4 + 64\left(\frac{\lambda}{R}\right)^4 \right\} \\ & + \frac{16}{21}(1+e)\nu g_0 R^4 \left\{ 32 - (2 + \cos 4\phi)\left(\frac{\eta}{R}\right)^2 + 12\left(\frac{\lambda}{R}\right)^2 \right\} = 0, \end{aligned} \quad (\text{C } 5)$$

$$\begin{aligned} & 35\sqrt{\pi}\eta R \cos 2\phi + (1+e)\nu g_0 [32(1+3e)R^2 - 3(3-e)(\eta^2 + 21\lambda^2) \\ & - 8\sqrt{\pi}(4-3e)\eta R \cos 2\phi] - \frac{210\sqrt{\pi}\lambda^2 R}{St_d} \\ & + \frac{16}{33}(1+e)\nu g_0 R^4 \left\{ 32(5+3e) + 2(2 + \cos 4\phi)(2-3e)\left(\frac{\eta}{R}\right)^2 \right. \\ & \left. - 33(5-3e)\left(\frac{\lambda}{R}\right)^2 - \frac{9}{32}(3-e)\left[\left(\frac{\eta}{R}\right)^4 - 11\left(\frac{\eta}{R}\right)^2\left(\frac{\lambda}{R}\right)^2 - 66\left(\frac{\lambda}{R}\right)^4\right] \right\} = 0, \end{aligned} \quad (\text{C } 6)$$

$$\begin{aligned} & 5\sqrt{\pi}R \cos 2\phi - (1+e)\nu g_0 [3(3-e)\eta + 2\sqrt{\pi}(1-3e)R \cos 2\phi] - \frac{10\sqrt{\pi}\eta R}{St_d} \\ & + 5\sqrt{\pi}\lambda^2 R \cos 2\phi - (1+e)\nu g_0 R^3 \left\{ \frac{8}{7}\sqrt{\pi}[(4-3e)\left(\frac{\lambda}{R}\right)^2 - 8(1+e)] \cos 2\phi \right. \\ & \left. + \frac{1}{42}\left(\frac{\eta}{R}\right)[64(4-3e) - 32(5+3e) \cos 4\phi - 3(3-e)\left(\frac{\eta}{R}\right)^2 + 36(3-e)\left(\frac{\lambda}{R}\right)^2] \right\} = 0, \end{aligned} \quad (\text{C } 7)$$

$$\begin{aligned} & 5(\eta - \sin 2\phi) + 2(1+e)(1-3e)\nu g_0 \sin 2\phi - 5\lambda^2 \sin 2\phi \\ & - \frac{8}{7}(1+e)\nu g_0 R^2 \sin 2\phi \left\{ 8(1+e) - (4-3e)\left(\frac{\lambda}{R}\right)^2 + \frac{4}{3\sqrt{\pi}}(5+3e)\frac{\eta}{R} \cos 2\phi \right\} = 0. \end{aligned} \quad (\text{C } 8)$$

Note that equations (C 5-C 6) contain only “even” order terms in $(\eta, \sin 2\phi, \lambda, R)$, with the neglected terms being of order six and beyond; on the other hand, equations (C 7-C 8) contain only “odd” order terms in $(\eta, \sin 2\phi, \lambda, R)$, with the neglected terms being of order five and beyond. Therefore, equations (C 5-C 8) belong to the “super-super-Burnett” orders since they incorporate all terms up-to “quartic-order” in the shear-rate ($R \sim \dot{\gamma}$).

C.1. Exact solution at Burnett order for the whole range of density

Removing the terms within the second-brackets in (C 5-C 8), we obtain the Burnett-order equations as given in (4.5). The latter equations admit an exact solution, given by (4.10a) and (4.11), as discussed in §4.1.

C.2. Perturbation solutions beyond Burnett order

The super-Burnett and super-super-Burnett order equations [(C 5-C 8)] are solved using a regular perturbation expansion around the exact Burnett-order solution (4.10a) and (4.11):

$$\left. \begin{aligned} \eta &= \eta^{(2)} + \varepsilon\eta^{(3)} + \varepsilon^2\eta^{(4)} \\ \lambda &= \lambda^{(2)} + \varepsilon\lambda^{(3)} + \varepsilon^2\lambda^{(4)} \\ R &= R^{(2)} + \varepsilon R^{(3)} + \varepsilon^2 R^{(4)} \\ \sin 2\phi &= \sin^{(2)} 2\phi + \varepsilon \sin 2\phi^{(3)} + \varepsilon^2 \sin 2\phi^{(4)} \end{aligned} \right\}. \quad (\text{C } 9)$$

In the above expressions $\varepsilon \sim \dot{\gamma}$ and the superscript “2” corresponds to the “Burnett-order” solution [i.e. the closed form expressions (4.10a) and (4.11) as given in §4.1] and the superscripts “3” and “4” correspond to the corrections at third and fourth orders, respectively.

Plugging (C 9) into corresponding third and fourth [(C 5-C 8)] order equations and

after performing some cumbersome algebra we obtain a null-solution at third-order

$$\eta^{(3)} = 0 = \lambda^{(3)} = R^{(3)} = \sin 2\phi^{(3)}. \quad (\text{C } 10)$$

The solutions at fourth-order are given by :

$$\begin{aligned} \eta^{(4)} = & - \left[\left[\sqrt{\pi}(1+e)\nu g_0 \cos 2\phi^{(2)} \{5 - 2(1+e)(1-3e)\nu g_0\} \left\{ 1024(5+3e)R^{(2)4} \right. \right. \right. \\ & - 192(1+3e)R^{(2)2} \left(\eta^{(2)2} - 4\lambda^{(2)2} \right) - 9(1-e) \left(\eta^{(2)4} - 8\eta^{(2)2}\lambda^{(2)2} + 84\lambda^{(2)4} \right) \left. \left. \right\} \right] \\ & - \left[8 \left\{ 5\sqrt{\pi}\eta^{(2)} \cos 2\phi^{(2)} + 2(1+e)\nu g_0 \left(8(1+3e)R^{(2)} - (1-3e)\sqrt{\pi}\eta^{(2)} \cos 2\phi^{(2)} \right) \right\} \right. \\ & \times \left\{ 210\sqrt{\pi}\lambda^{(2)2} R^{(2)} \cos 2\phi^{(2)} - 48(1+e)\sqrt{\pi}\nu g_0 R^{(2)} \cos 2\phi^{(2)} \left((4-3e)\lambda^{(2)2} - 8(1+e)R^{(2)2} \right) \right. \\ & \left. \left. - 3(1+e)\nu g_0 \eta^{(2)} \left(32(1-3e)R^{(2)2} - (3-e)(\eta^{(2)2} - \lambda^{(2)2}) \right) \right\} \right] \right] \\ & \frac{168 \left[\sqrt{\pi} \cos 2\phi^{(2)} \left\{ 2\sqrt{\pi} \cos 2\phi^{(2)} \{5 - 2(1+e)(1-3e)\nu g_0\} R^{(2)} - 3(1-e^2)\nu g_0 \eta^{(2)} \right\} \right.}{168 \left[\sqrt{\pi} \cos 2\phi^{(2)} \left\{ 2\sqrt{\pi} \cos 2\phi^{(2)} \{5 - 2(1+e)(1-3e)\nu g_0\} R^{(2)} - 3(1-e^2)\nu g_0 \eta^{(2)} \right\} \right.} \\ & \times \{5 - 2(1+e)(1-3e)\nu g_0\} + 2\nu g_0 \left\{ 3(1+e)(3-e) + 10\sqrt{\pi}R_{St}^{(2)} \right\} \left\{ 5\sqrt{\pi}\eta^{(2)} \cos 2\phi^{(2)} \right. \\ & \left. \left. + 2(1+e)\nu g_0 \left(8(1+3e)R^{(2)} - (1-3e)\sqrt{\pi}\eta^{(2)} \cos 2\phi^{(2)} \right) \right\} \right] \quad (\text{C } 11) \end{aligned}$$

$$\begin{aligned} \lambda^{(4)} = & \frac{1}{77616\lambda^{(2)}} \left(\left[\frac{28(1+e)}{\left\{ 3(1+e)(3-e) + 10\sqrt{\pi}R_{St}^{(2)} \right\}} \left\{ 1024(5+3e)R^{(2)4} \right. \right. \right. \\ & + 96R^{(2)2} \left(2(2-3e)\eta^{(2)2} - 11(5-3e)\lambda^{(2)2} \right) - 9(3-e) \left(\eta^{(2)4} - 11\eta^{(2)2}\lambda^{(2)2} - 66\lambda^{(2)4} \right) \left. \left. \right\} \right]_a \\ & - \left[\frac{132}{\left\{ 3(1+e)(3-e) + 10\sqrt{\pi}R_{St}^{(2)} \right\} \sqrt{\pi} \{5 - 2(1+e)(1-3e)\nu g_0\} \nu^2 g_0^2 \cos 2\phi^{(2)}} \right. \\ & \times \left[35\sqrt{\pi}\eta^{(2)} \cos 2\phi^{(2)} + 8(1+e)\nu g_0 \left\{ 8(1+3e)R^{(2)} - (4-3e)\sqrt{\pi}\eta^{(2)} \cos 2\phi^{(2)} \right\} \right] \\ & \times \left[70\sqrt{\pi}\lambda^{(2)2} R^{(2)} \cos 2\phi^{(2)} + (1+e)\nu g_0 \left\{ 16\sqrt{\pi}R^{(2)} \cos 2\phi^{(2)} \left(8(1+e)R^{(2)2} - (4-3e)\lambda^{(2)2} \right) \right. \right. \\ & \left. \left. - 32(1-3e)\eta^{(2)2} R^{(2)} + (3-e)\eta^{(2)} \left(\eta^{(2)2} - 12\lambda^{(2)2} \right) \right\} \right] \right]_b \\ & + \left[\frac{1848\eta^{(4)} \left[\frac{\sqrt{\pi} \{35 - 8(1+e)(4-3e)\nu g_0\} R^{(2)} \cos 2\phi^{(2)} - 6(3-e)(1+e)\nu g_0 \eta^{(2)}}{\left\{ 3(1+e)(3-e)\nu g_0 + 10\sqrt{\pi}\nu g_0 R_{St}^{(2)} \right\}} \right. \right. \\ & \left. \left. + \frac{\left\{ 35\sqrt{\pi}\eta^{(2)} \cos 2\phi^{(2)} + 8(1+e)\nu g_0 \left(8(1+3e)R^{(2)} - (4-3e)\sqrt{\pi}\eta^{(2)} \cos 2\phi^{(2)} \right) \right\}}{\sqrt{\pi} \{5 - 2(1+e)(1-3e)\nu g_0\} \cos 2\phi^{(2)}} \right] \right]_c \quad (\text{C } 12) \end{aligned}$$

$$\begin{aligned}
R^{(4)} = \frac{1}{42\sqrt{\pi}\{5 - 2(1+e)(1-3e)\nu g_0\} \cos 2\phi^{(2)}} & \left[-210\sqrt{\pi}\lambda^{(2)^2} R^{(2)} \cos^{(2)} 2\phi \right. \\
& + 48(1+e)\sqrt{\pi}\nu g_0 R^{(2)} \cos^{(2)} 2\phi \left\{ (4-3e)\lambda^{(2)^2} - 8(1+e)R^{(2)^2} \right\} \\
& + 3(1+e)\nu g_0 \eta^{(2)} \left\{ 32(1-3e)R^{(2)^2} - (3-e)(\eta^{(2)^2} - 12\lambda^{(2)^2}) \right\} \\
& \left. + 42\eta^{(4)} \left\{ 3(3-e)(1+e) + 10\sqrt{\pi}R_{St}^{(2)} \right\} \nu g_0 \right] \quad (C 13)
\end{aligned}$$

$$\begin{aligned}
\sin 2\phi^{(4)} = \frac{1}{21\sqrt{\pi}\{5 - 2(1+e)(1-3e)\nu g_0\}} & \left[105\sqrt{\pi}(\eta^{(4)} - \lambda^{(2)^2} \sin^{(2)} 2\phi) \right. \\
& \left. - 8(1+e)\nu g_0 \sin^{(2)} 2\phi \left\{ 4(5+3e)\eta^{(2)} R^{(2)} \cos^{(2)} 2\phi - 3\sqrt{\pi}((4-3e)\lambda^{(2)^2} - 8(1+e)R^{(2)^2}) \right\} \right] \quad (C 14)
\end{aligned}$$

where $R_{St}^{(2)} = R^{(2)}/\nu g_0 St_d$. In absence of interstitial gas, i.e., in the limit of $St_d \rightarrow \infty$, the St -dependent terms in (C 11-C 14) disappear and we obtain the corresponding super-super-Burnett order solutions for the dry-granular flow – the resulting expressions match exactly with the solution provided by Saha & Alam (2016) for the uniform shear flow of dry granular fluid.

In the dilute limit, the solutions at fourth-order are given by:

$$\begin{aligned}
\eta^{(4)} = \frac{1}{56\{3(5-e)(1+e)\eta^{(2)} + 10\sqrt{\pi}R_{St}^{(2)}(St \cos 2\phi + 2\eta^{(2)})\}} & \left\{ (1+e)(27-11e)\eta^{(2)^4} \right. \\
& \left. + 252(1-e^2)\lambda^{(2)^4} + 560\sqrt{\pi} \cos 2\phi \eta^{(2)} \lambda^{(2)^2} St R_{St}^{(2)} - 24(1+e)(13-5e)\eta^{(2)^2} \lambda^{(2)^2} \right\}, \quad (C 15)
\end{aligned}$$

$$\begin{aligned}
\lambda^{(4)} = \frac{7(1+e)}{17248\lambda^{(2)}\{3(3-e)(1+e) + 10\sqrt{\pi}R_{St}^{(2)}\}} & \left\{ 132(19-11e)\lambda^{(2)^4} \right. \\
& \left. - (13+3e)\eta^{(2)^4} + 176\eta^{(2)^2}\lambda^{(2)^2} - 88(5+3e)\eta^{(2)}\eta^{(4)} \right\}, \quad (C 16)
\end{aligned}$$

$$\begin{aligned}
R^{(4)} = \frac{\nu}{560\sqrt{\pi}\eta^{(2)} \cos^{(2)} 2\phi} & \left\{ 3(1-e^2)(\eta^{(2)^4} - 8\eta^{(2)^2}\lambda^{(2)^2} + 84\lambda^{(2)^4} + 56\eta^{(2)}\eta^{(4)}) \right. \\
& \left. - 560\sqrt{\pi} \cos 2\phi St R_{St}^{(2)} \eta^{(4)} \right\}, \quad (C 17)
\end{aligned}$$

$$\sin 2\phi^{(4)} = \left\{ \eta^{(4)} - \lambda^{(2)^2} \sin^{(2)} 2\phi \right\}. \quad (C 18)$$

where $R_{St}^{(2)} = R^{(2)}/\nu$.

C.3. Burnett-order equations for dilute gas-solid suspension: scaling arguments

Considering the dry granular limit ($St \rightarrow \infty$) of a dilute gas-solid suspension, the Burnett-order second-moment balance equations are given by:

$$\left. \begin{aligned} 20\sqrt{\pi}\eta R \cos 2\phi - 3(1-e^2)\nu(10+\eta^2) &= 0 \\ 35\sqrt{\pi}\eta R \cos 2\phi - 3(1+e)(3-e)\nu(\eta^2 + 21\lambda^2) &= 0 \\ 5\sqrt{\pi}R \cos 2\phi - 3(1+e)(3-e)\nu\eta &= 0 \\ \eta - \sin 2\phi &= 0 \end{aligned} \right\}. \quad (C 19)$$

This admits an analytical solution of the form (Saha & Alam 2016)

$$\eta^2 = \frac{10(1-e)}{(11-3e)}, \quad \lambda^2 = \frac{20(1-e)}{7(11-3e)}, \quad \sin 2\phi = \eta, \quad R = \frac{3(3-e)(1+e)}{5\sqrt{\pi} \cos 2\phi} \frac{\sqrt{10}}{\sqrt{11-3e}} \nu \sqrt{1-e} \quad (\text{C } 20)$$

At leading order, we have

$$\eta = \frac{\sqrt{5(1-e)}}{2}, \quad \lambda = \sqrt{\frac{5(1-e)}{14}}, \quad \sin 2\phi = \frac{\sqrt{5(1-e)}}{4}, \quad R = \frac{6}{\sqrt{5\pi}} \nu \sqrt{1-e}, \quad (\text{C } 21)$$

as was predicted also by Richman (1989). It is easy to verify that the following quantities,

$$\eta, \lambda, \sin 2\phi, R/\nu, \sim \sqrt{1-e}, \quad (\text{C } 22)$$

are of the same order which holds for a sheared dilute granular gas. Equation (C 22) helps to prove a scaling between the Stokes number and the inelasticity.

Proposition: The following scaling,

$$St \sim \sqrt{1-e}, \quad (\text{C } 23)$$

holds for a sheared dilute gas-solid suspension.

To clarify (C 23), let us consider the second-moment balance equations for a dilute suspension at the Burnett-order:

$$20\sqrt{\pi}\eta R \cos 2\phi - 3(1-e^2)\nu(10+\eta^2) - \frac{60\sqrt{\pi}R}{St} = 0 \quad (\text{C } 24a)$$

$$35\sqrt{\pi}\eta R \cos 2\phi + (1+e)\nu[-3(3-e)(\eta^2+21\lambda^2)] - \frac{210\sqrt{\pi}\lambda^2 R}{St} = 0, \quad (\text{C } 24b)$$

$$5\sqrt{\pi}R \cos 2\phi - 3(3-e)(1+e)\nu\eta - \frac{10\sqrt{\pi}\eta R}{St} = 0, \quad (\text{C } 24c)$$

$$5(\eta - \sin 2\phi) = 0. \quad (\text{C } 24d)$$

From (C 24c), we find

$$5\sqrt{\pi}R \cos 2\phi \sim R \sim \nu\sqrt{1-e} \quad \text{and} \quad 3(1+e)(3-e)\nu\eta \sim \nu\sqrt{1-e},$$

and hence

$$\frac{10\sqrt{\pi}\eta R}{St} \sim \frac{\sqrt{1-e}\nu\sqrt{1-e}}{\sqrt{1-e}} \sim \nu\sqrt{1-e},$$

which implies that all three terms in (C 24c) are of the same order. Similarly, from (C 24b) we have

$$35\sqrt{\pi}\eta R \cos 2\phi \sim \nu(1-e) \quad \text{and} \quad 3(1+e)(3-e)\nu(\eta^2+21\lambda^2) \sim \nu(1-e),$$

and hence

$$\frac{210\sqrt{\pi}\lambda^2 R}{St} \sim \frac{(1-e)\nu\sqrt{1-e}}{\sqrt{1-e}} \sim \nu(1-e),$$

again implying that three terms in (C 24b) are of the same order. From the above ordering arguments the proposition made in (C 23) is justified.

Acknowledgements

This work is funded by the Department of Science and Technology under “India-Netherlands” project (Ref.: DST/INT/NL/P-03/2016); S.S. was supported by this project via a postdoctoral fellowship. This work was conceived, supervised and written by M.A.

REFERENCES

- ALAM, M. & SAHA, S. 2017 Normal stress differences and beyond-Navier-Stokes hydrodynamics. *EPJ Conf. Proc.* **140**, 11014.
- ALAM, M., SAHA, S. & GUPTA, R. 2019 Unified theory for a sheared gas-solid suspension: from rapid-granular suspension to its small Stokes-number limit. *J. Fluid Mech.*, **870**, 1175-1193.
- RICHMAN, M. W. 1989 The source of second moment in dilute granular flows of highly inelastic spheres. *J. Rheol.* **33**, 1293-1306.
- SAHA, S. & ALAM, M. 2014 Non-Newtonian stress, collisional dissipation and heat flux in the shear flow of inelastic disks: a reduction via Grad's moment method. *J. Fluid Mech.* **757**, 251-296.
- SAHA, S. & ALAM, M. 2016 Normal stress differences, their origin and constitutive relations for a sheared granular fluid. *J. Fluid Mech.* **795**, 549-580.
- SAHA, S. & ALAM, M. 2017 Revisiting ignited-quenched transition and the non-Newtonian rheology of a sheared dilute gas-solid suspension. *J. Fluid Mech.* **833**, 206-246.
- SAHA, S. & ALAM, M. 2020 Burnett-order constitutive relations, second moment anisotropy and co-existing states in sheared dense gas-solid suspensions. *J. Fluid Mech.*, **887**, art. no. ?? (In press)