

Supplementary material for Asymptotics of stream-wise Reynolds stress in wall turbulence

by Peter A. Monkewitz (peter.monkewitz@epfl.ch)

1 Alternative asymptotic sequences for the inner expansion of $\langle uu \rangle^+$

1.1 First alternative sequence $[1, 1/\ln(\text{Re}_\tau), \dots]$

The first alternative asymptotic expansion of the near-wall stream-wise normal stress,

$$\langle uu \rangle_{\text{DNS}}^+ = \tilde{f}(y^+) + \tilde{g}(y^+)/\ln(\text{Re}_\tau) \quad (1)$$

corresponds essentially to the one proposed by Monkewitz & Nagib (2015). Using the DNS profiles in table 1 of the main paper to determine the two terms of the asymptotic expansion (1) results in figure 1 of this supplement. The figure shows, that the different DNS pairs yield consistent results for $\tilde{f}(y^+)$ and $\tilde{g}(y^+)$ between the wall and $y^+ \approx 200$, similar to figure 1 of the main paper.

The expansion (1) also yields an inner peak height of

$$\left[\tilde{f} + \tilde{g}/\ln(\text{Re}_\tau) \right]_{\text{IP}} = 14.0 - 40.9/\ln(\text{Re}_\tau) \quad (2)$$

which is in excellent agreement with figure 2 of the main paper. However, the resulting coefficient of $(y^+)^2$ in the Taylor expansion of $\langle uu \rangle^+$ about $y^+ = 0$ is $(0.30 - 0.86/\ln \text{Re}_\tau)$, i.e. significantly overshoots the maximum of $1/4$ posited by Chen & Sreenivasan (2021) for $\text{Re}_\tau \rightarrow \infty$. Therefore, the inner expansion (1) is not further pursued here.

1.2 Second alternative sequence $[\ln(\text{Re}_\tau), 1, \dots]$

It is instructive to test the asymptotic sequence $[\ln(\text{Re}_\tau), 1, \dots]$ of the attached eddy model (Marusic & Monty, 2019) with the same methodology, i.e. decompose the channel DNS profiles of table 1 in the main paper according to

$$\langle uu \rangle_{\text{DNS}}^+ = \tilde{f}(y^+) + \tilde{g}(y^+) \ln(\text{Re}_\tau) \quad (3)$$




Figure 1: The $\mathcal{O}(1)$ (panel a) and $\mathcal{O}(1/\ln(\text{Re}_\tau))$ (panel b) components of $\langle uu \rangle_{\text{wall}}^+$, extracted from the same channel DNS pairs of table 1 in the main paper, using the same color scheme as in its figure 1: (red) profiles 1 and 2 (—), 1 and 3 (- - -), 1 and 4 (\cdots), 1 and 5 ($- \cdot -$); (green) 2 and 3 (—), 2 and 4 (- - -), 2 and 5 (\cdots); (blue) 3 and 5 (—), 4 and 5 (- - -). \bullet , inner peak (equation 2 of this supplement); — and $- \cdot -$, fits similar to equation (2.2) of main paper.

The result is shown in figure 2. While the resulting inner peak at $y^+ \cong 15$ is quite well fitted by

$$\left[\tilde{f} + \tilde{g} \ln(\text{Re}_\tau) \right]_{\text{IP}} = 3.54 + 0.646 \ln(\text{Re}_\tau) \quad , \quad (4)$$

given as equation (3.2) in Samie *et al.* (2018), the \tilde{f} 's and \tilde{g} 's from different profile pairs are seen to spread out, indicating that the asymptotic sequence $[\ln(\text{Re}_\tau), 1, \dots]$ is not appropriate (see also the formal argument against this scaling in the main text).




Figure 2: The $\mathcal{O}(\ln(\text{Re}_\tau))$ (panel a) and $\mathcal{O}(1)$ (panel b) components of $\langle uu \rangle^+$, extracted from the same channel DNS pairs of table 1 in the main paper, using the same color scheme as in its figure 1. \bullet , inner peak of Samie *et al.* (2018) (equation (4) of this supplement).

2 Comparison of the present MAE with the fit of Marusic and Kunkel for $\langle uu \rangle^+$

The composite MAE profiles of figure 3 of the main paper are compared to the fits of Marusic & Kunkel (2003), inspired by the attached eddy model (see also Monkewitz *et al.*, 2017). Note, that the inner and outer parts of these fits are **not matched across an overlap region**, but patched across the interval $30 \leq y^+ \leq 150$ with a cubic in $\ln(y^+)$.

All the parameters are as in Marusic & Kunkel (2003) and Marusic & Perry (1997), except for

$$(A1, B1) = (1.26, 2.1) \quad , \quad (5)$$

instead of the original (1.03, 2.39). This value of $A1$ has been proposed by Marusic *et al.* (2013) and the value of $B1$, which is well within the range of their table 2, has been chosen to provide a good fit to the outer part of the ZPG TBL profile of Samie *et al.* (2018) at $\text{Re}_\tau = 20000$. Beyond this, no attempt has been made to modify the parameters of Marusic & Kunkel (2003), but simultaneously matching all the high quality profiles included in figure 3 appears difficult.




Figure 3: Solid lines: composite profiles of $\langle uu \rangle^+$ as in figure 3 of the main paper for $Re_\tau = 1000$ (green), 1995 (blue), 5186 (red), 20'250 (green), 98'190 (blue), 10^6 (red), 10^7 (green), 10^9 (blue). - - - (black) corresponding fits of Marusic & Kunkel (2003). DNS and experimental profiles: - - -, DNS profiles 5 (green), 4 (blue) and 1 (red) of table 1 in the main paper. $\triangle\triangle\triangle$, NSTAP Superpipe profiles of Hultmark *et al.* (2012) at $Re_\tau = 1985, 5411, 20'250$ and 98'187. \square , ZPG TBL profile of Sillero *et al.* (2013) for $Re_\tau = 1989$ (blue) and of Samie *et al.* (2018) for $Re_\tau = 20'000$ (orange).

References

- CHEN, XI & SREENIVASAN, KATEPALLI R. 2021 Reynolds number scaling of the peak turbulence intensity in wall flows. *Journal of Fluid Mechanics* **908**, R3.
- HULTMARK, M., VALLIKIVI, M., BAILEY, S. C. & SMITS, A. J. 2012 Turbulent pipe flow at extreme Reynolds numbers. *Phys. Rev. Lett.* **108**, 094501–1–5.
- MARUSIC, I. & KUNKEL, G. J. 2003 Streamwise turbulence intensity formulation for flat-plate boundary layers. *Phys. Fluids* **15**, 2461–2464.
- MARUSIC, IVAN & MONTY, JASON P. 2019 Attached eddy model of wall turbulence. *Annual Review of Fluid Mechanics* **51** (1), 49–74.
- MARUSIC, I., MONTY, J. P., HULTMARK, M. & SMITS, A. J. 2013 On the logarithmic region in wall turbulence. *J. Fluid Mech. Rapids* **716**, R3.
- MARUSIC, I., UDDIN A.K.M. & PERRY, A.E. 1997 Similarity law for the streamwise turbulence intensity in zero-pressure-gradient turbulent boundary layers. *Physics of Fluids* **9**, 3718–3726.
- MONKEWITZ, P. A. & NAGIB, H. M. 2015 Large Reynolds number asymptotics of the stream-wise normal stress in ZPG turbulent boundary layers. *J. Fluid Mech.* **783**, 474–503.
- MONKEWITZ, P. A., NAGIB, H. M. & BOULANGER, V. 2017 Comparing the three possible scalings of stream-wise normal stress in turbulent boundary layers. In *Proc. TSFP10, Chicago, (USA)* (ed. A. J. Smits & H. M. Nagib). www.tsfp-conference.org.
- SAMIE, M., MARUSIC, I., HUTCHINS, N., FU, M. K., FAN, Y., HULTMARK, M. & SMITS, A. J. 2018 Fully resolved measurements of turbulent boundary layer flows up to $Re_\tau=20\,000$. *Journal of Fluid Mechanics* **851**, 391–415.
- SILLERO, JUAN A., JIMÉNEZ, JAVIER & MOSER, ROBERT D. 2013 One-point statistics for turbulent wall-bounded flows at Reynolds numbers up to $\delta^+ \approx 2000$. *Physics of Fluids* **25**, 105102.