

**Life-history traits of the striped red mullet *Mullus surmuletus* (Linnaeus, 1758)  
in the south Aegean Sea (eastern Mediterranean)**

**Vasiliki Kousteni<sup>\*1</sup>, Aikaterini Anastasopoulou and Chryssi Mytilineou**

Institute of Marine Biological Resources and Inland Waters, Hellenic Centre for Marine Research,  
46.7<sup>th</sup> km Athens Sounio ave., P.O. Box 712, 19013 Anavyssos Attiki, Greece

**Supplementary Material**

---

<sup>\*1</sup>Corresponding author: Vasiliki Kousteni

Institute of Marine Biological Resources and Inland Waters, Hellenic Centre for Marine Research, 46.7<sup>th</sup> km Athens Sounio ave., P.O. Box 712, 19013 Anavyssos Attiki, Greece. Email: kousteni@hcmr.gr. Tel.: +30 210 9856714.

**Table S1.** Length at 50% maturity ( $L_{50}$ ), VBGF parameters ( $L_{\text{inf}}$ ,  $k$ ,  $t_0$ ), maximum observed age ( $A_m$ ), size range ( $L_T$ ) and length-weight relationship parameters ( $a$ ,  $b$ ,  $r^2$ ) of *Mullus surmuletus* in different locations (N, number of individuals; F, female; M, male; T, combined sexes;  $L_F$ , fork length).

Location	Sex	N	$L_{50}$	N	$L_T$ range	$\alpha$	$b$	$r^2$	N	$L_{\text{inf}}$	$k$	$t_0$	$A_m$	Method	Reference
Eastern Mediterranean Sea															
South Aegean Sea (Greece)	F	593	153.3	593	103–320 $L_T$	0.0155	2.915	0.96	593	346.1	0.299	-0.984	5	otolith	<b>Present study</b>
	M	279	139.2	279	121–162 $L_T$	0.0032	2.976	0.98					3		
	T		1032	95–320 $L_T$	0.0013	2.943	0.96		872	373.2	0.255	-0.999	5		
Central Aegean Sea (Greece)	F	336	138.4	336	90–260 $L_F$	6E-06	3.229	0.99	336	413.0	0.100	-2.800	6	otolith	Vassilopoulou & Papaconstantinou (1992)
	M	451	115.5	451	100–200 $L_F$	5E-06	3.254	0.97		380.0	0.100	-2.760	5		
South Evoikos Gulf (Greece)	T		307	101–201 $L_F$	9E-06	3.140	0.97								Petrakis & Stergiou (1995)
Cyclades (Off Naxos, Greece)	T		257	138–320 $L_T$	0.0140	2.954	0.94								Moutopoulos & Stergiou (2002)
Off Crete Island (Greece)	T								1526	354.0	0.225	-1.194	5	scale	Machias <i>et al.</i> (1998)
Thermaikos Gulf and Thrakian Sea (Greece)	F								248.0	0.260	-1.580			otolith	Papaconstantinou <i>et al.</i> (1994)
	M								220.0	0.270	-1.460				
	T		292	70–240 $L_F$	0.0116	3.149	0.99			0.270	-1.460				
NE Aegean Sea (Greece)	T		48	44–97 $L_T$	0.0045	3.510	0.99								Koutrakis & Tsikliras (2003)
Marmara Sea (Turkey)	F								344.8	0.210	-2.970				Moldur (1999)
	M								273.0	0.250	-2.110				
	T								328.3	0.230	-2.130				
North Aegean Sea (Turkey)	T		601	109–299 $L_T$	0.0069	3.192	0.98								Karakulak <i>et al.</i> (2006)
Saros Bay (eastern Aegean Sea, Turkey)	F	132.0	184	110–268	0.0110	3.020	0.95		124	283.8	0.190	-2.160	7	otolith	Arslan & İşmen (2013)
	M	137.0	118	118–198	0.0070	3.162	0.97		89	269.4	0.200	-2.340	3		
	T	656	656	96–268					272	278.2	0.200	-2.160	7		
Edremit Bay (eastern Aegean Sea, Turkey)	T	250.0	520	77–170 $L_T$	0.0044	3.356	0.94								Torcu-Koç <i>et al.</i> (2015)
	T								250.9	0.140	-2.480				Üstün (2010)
Izmir Bay (eastern Aegean Sea, Turkey)	T		192	66–226 $L_T$	0.0083	3.127	0.98		278.5	0.193	-1.578			otolith	İlhan <i>et al.</i> (2009)
	T		38	73–186 $L_T$	0.0131	2.970	0.98								Gökçe <i>et al.</i> (2007)
	T		117	74–619 $L_T$	0.0106	3.202	0.99								Özaydin & Taşkavak (2007)
Gulf of Antalya (Turkey)	T		45	137–245 $L_T$	0.0029	3.465	0.95								Özvarol (2014)
Babadillimanı Bight (Turkey)	T		145	55–222 $L_T$	0.0082	3.110	0.98								Çicek <i>et al.</i> (2006)
Off Egypt	F	150.0													Hashem (1973)
	M	130.0													
	T		122	54–208 $L_T$	0.0110	3.030	0.86								Abdallah (2002)
	T	1835	151.0	1835	50–291 $L_T$	0.0104	3.062	0.98	1835	317.4	0.470	-0.300	5	otolith	Mehanna (2009)

Table S1 (continue)

Location	Sex	N	$L_{50}$	N	$L_T$ range	$\alpha$	b	$r^2$	N	$L_{inf}$	k	$t_0$	$A_m$	Method	Reference
Central Mediterranean Sea															
Southern Tyrrhenian and Ionian Seas (Italy)	F								301.2	0.240	-2.390		otolith	Andaloro (1982)	
	M								250.2	0.300	-2.680				
	T			40–280	$L_T$	0.0067	3.205		311.0	0.240	-2.680				
Eastern Adriatic (Croatia)	T	127	154–309	$L_T$	3E-07	3.512	0.89								Dulčić & Kraljević (1996)
	T	47	125–285	$L_T$	0.0039	3.367	0.96								Dulčić & Glamuzina (2006)
Strait of Sicily (Italy)	F								297.5	0.490	-0.310		otolith	Andaloro & Giarritta (1985)	
	M								262.5	0.410	-0.230				
	T		98–267		0.0093	3.070									
	T				0.0093	3.000									Djabali <i>et al.</i> (1993)
Off Tunisia	F	155.0			0.1443	3.280			218.2	0.510	-0.112		scale	Gharbi & Ktari (1981)	
	M	130.0			0.1403	3.351			198.7	0.490	-0.025				
	T		80–230						202	215.1	0.500	-0.116	5		
	F								212.0	0.430	-0.650		otolith	Jabeur (1999)	
	M								226.0	0.280	-1.070				
	T	123	50–230						123	223.0	0.344	-0.790	6		
Western Mediterranean Sea															
Off Morocco	F								323.0	0.540	-0.620		otolith	Lamrini (2010)	
	M								303.0	0.520	-0.160				
	F	1037	123–381	$L_T$	0.0072	3.178	0.98		394.0	0.190	-1.030		otolith	Bakali <i>et al.</i> (2016)	
	M	904	112–345	$L_T$	0.0068	3.173	0.97		356.4	0.130	-2.170				
	T	2033	2033	112–381	$L_T$	0.0071	3.173	0.98	1520	392.0	0.130	-3.210	10		

Table S1 (continue)

Location	Sex	N	$L_{50}$	N	$L_T$ range	$\alpha$	b	$r^2$	N	$L_{inf}$	k	$t_0$	$A_m$	Method	Reference	
Western Mediterranean Sea																
Off Balearics	F								345.3	0.137	-3.821			otolith	Morales-Nin (1991)	
	M								232.9	0.288	-3.325					
	T			95–270	$L_T$	0.0160	2.913		297.6	0.238	-2.640					
	T	155.0		95–270	$L_T$				298.0	0.240	-2.320	4			Morales-Nin (1992)	
	F	604	168.0	1866	120–320	$L_T$	0.0095	3.109	0.97	1771	319.0	0.205	-2.605	6	otolith	Reñones <i>et al.</i> (1995)
	M	662	150.0	1429	100–280	$L_T$	0.0105	3.067	0.97	1342	255.4	0.273	-2.450	6		
	T			3541	100–320	$L_T$	0.0091	3.120	0.98	3339	312.8	0.211	-2.348	6		
	F	322	177.0	322	120–240	$L_T$	0.0090	3.020		322	247.0	0.370	-0.370		LFD	Kherraz <i>et al.</i> (2014)
	M	516		516	125–235	$L_T$	0.0100	2.980		516	255.2	0.320	-0.710			
Off Algeria (Mostaganem coast)	T	838		838	120–235	$L_T$	0.0090	2.934		838				8		
	T				120–320		0.0073	3.100		3339	325.2	0.110	-3.647	6	otolith	Sánchez <i>et al.</i> (1983)
	T				50–200				309.4	0.110	-3.850			otolith	Morales-Nin (1986)	
Catalan Sea (Spain)	F								334.0	0.430	-0.600			NA	Campillo (1992)	
	M								284.0	0.530	-0.440					
	T				$L_T$	0.0182	3.000								Djabali <i>et al.</i> (1993)	
	T					0.0067	3.000									
Eastern Atlantic Ocean																
Bay of Biscay (Spain)	T	493	155.0	493	166–436				493	377.0	0.290		9	otolith	N'Da (1992)	
	F								427.0	0.280	0.641				N'Da <i>et al.</i> (2006)	
	M								359.0	0.300	0.740					
Eastern English Channel and southern North Sea	F	169.0	736	114–425	$L_T$	3.2E-06	3.352	0.95	551	512.0	0.196	-2.904	7	otolith	Mahé <i>et al.</i> (2013)	
	M	162.0	353	77–326	$L_T$	4.8E-06	3.177	0.98	320	364.0	0.218	-3.232	6			
	T		1089	77–425	$L_T$	3.3E-06	3.243	0.97	383	475.0	0.210		7			
North Sea	T								533.4	0.180	-1.230				Mahé <i>et al.</i> (2005)	
Gulf of Cadiz (SW Spain)	T	466		94–354	$L_T$	0.0054	3.282	0.98							Torres <i>et al.</i> (2012)	
East coast of Spain	T	146		77–254	$L_S$	0.0097	3.075	1.00							Vale <i>et al.</i> (2003)	
Off Portugal (SW coast)	T	299		215–380	$L_T$	2.4E-06	3.087	0.92							Gonçalves <i>et al.</i> (1996)	
Off Canary Islands	F	169.0	541	140–330	$L_T$	0.0074	3.179	0.96	221	369.3	0.190	-1.980		otolith	Pajuelo <i>et al.</i> (1997)	
	M	163.0	235	140–260	$L_T$	0.0128	2.990	0.99	210	291.9	0.230	-1.760				
	T	166.0	1215	120–330	$L_T$	0.0074	3.183	0.99	723	357.1	0.220	-1.840	8			

## References (cited only in the SM)

- Abdallah M (2002) Length-mass relationship of fishes caught by trawl off Alexandria, Egypt. *NAGA, the ICLARM Quarterly* 25, 19–20.
- Çiçek E, Avşar D, Yeldan H and Özütok M (2006) Length-weight relationships for 31 teleost fish caught by bottom trawl net in the Babadillimani Bight (Northeastern Mediterranean). *Journal of Applied Ichthyology* 22, 290–292.
- Djabali F, Meħaili A, Koudil M and Brahmi B (1993) Empirical equations for the estimation of natural mortality in Mediterranean teleosts. *NAGA, the ICLARM Quarterly* 16, 35–37.
- Dulčić J and Glamuzina B (2006) Length-weight relationships for selected fish species from three eastern Adriatic estuarine systems (Croatia). *Journal of Applied Ichthyology* 22, 254–256.
- Dulčić J and Kraljević M (1996) Weight-length relationships for 40 fish species in the eastern Adriatic (Croatian waters). *Fisheries Research* 28, 243–251.
- Gökçe G, Aydin I and Metin C (2007) Length-weight relationships of 7 fish species from the North Aegean Sea, Turkey. *International Journal of Natural and Engineering Sciences* 1, 51–52.
- Gonçalves JMS, Bentes L, Lino PG, Ribeiro J, Canário AVM and Erzini K (1996) Weight-length relationships for selected fish species of the small scale demersal fisheries of the south and southwest coast of Portugal. *Fisheries Research* 30, 253–256.
- Karakulak FS, Erk H and Bilgin B (2006) Length-weight relationships for 47 coastal species from the Northern Aegean Sea, Turkey. *Journal of Applied Ichthyology* 22, 274–278.
- Koutrakis M and Tsikliras A (2003) Length-weight relationship of fishes from three northern Aegean estuarine systems (Greece). *Journal of Applied Ichthyology* 19, 258–260.
- Moutopoulos D and Stergiou KI (2002) Length-weight and length-length relationships of fish species from the Aegean Sea, Greece. *Journal of Applied Ichthyology* 17, 1–4.
- Özaydın O and Taşkavak E (2007) Length-weight relationships for 47 fish species from Izmir Bay (eastern Aegean Sea, Turkey). *Acta Adriatica* 47, 211–216.
- Özvarol Y (2014) Length-weight relationships of 14 fish species from the Gulf of Antalya (northeastern Mediterranean Sea, Turkey). *Turkish Journal of Zoology* 38, 342–346.

Petrakis G and Stergiou KI (1995) Weight-length relationships for 33 fish species in Greek waters. *Fisheries Research* 21, 465–469.

Torres MA, Ramos F and Sobrino I (2012) Length-weight relationships of 76 fish species from the Gulf of Cadiz (SW Spain). *Fisheries Research* 127–128, 171–175.

Vale C, Bayle JT and Ramos AA (2003) Weight and length relationships for selected fish species of the Western Mediterranean Sea. *Journal of Applied Ichthyology* 19, 261–262.