

Remnants of native forests support carnivore diversity in vineyard landscapes in central Chile

CAMILA B. GARCÍA, GABRIELLA L. SVENSSON, CAMILA BRAVO, MARÍA I. UNDURRAGA, JAVIERA DÍAZ-FORESTIER, KARINA GODOY
ALEXANDER NEAMAN, OLGA BARBOSA, SEBASTIÁN ABADES and JUAN L. CELIS-DIEZ

SUPPLEMENTARY TABLE 1 Number of carnivore detections in vineyard landscapes of central Chile (Fig. 1).

Site	<i>Lycalopex culpaeus</i>	<i>Lycalopex griseus</i>	<i>Canis familiaris</i>	<i>Felis catus</i>	<i>Leopardus guigna</i>	<i>Leopardus colocolo</i>	<i>Conepatus chinga</i>	<i>Galictis cuja</i>
1	50	4	7	0	0	1	0	2
2	6	0	0	0	0	0	0	0
3	21	0	5	1	0	0	0	0
4	1	40	0	0	29	0	7	4
5	0	31	0	0	1	0	0	0
6	1	4	17	0	1	0	0	0
7	1	0	24	2	7	0	0	1
8	7	12	1	0	9	0	0	0
9	11	0	20	0	0	0	0	0
10	0	17	0	0	4	0	0	3
11	0	12	4	0	10	0	0	0
12	0	15	5	1	7	0	0	0