

First insights into rodent trapping in Oku village, north-west Cameroon, based on interviews with local hunters

ALAIN D. MISSOUP, NGARTOUBAM DIDIER, ERNEST C. KEMING and CHRISTIANE DENYS

SUPPLEMENTARY TABLE 1 Mean weight and range (in g; Monadjem et al., 2015) for each of the 15 rodent taxa targeted by local hunters in Oku village, Cameroon.

Taxon	Mean weight (range) (g)
<i>Lophuromys</i>	63.7 (34–87)
<i>Praomys hartwigi</i>	57.2 (44–70)
<i>Hylomyscus</i> spp.	27.3 (16–41)
<i>Mastomys</i> spp.	63.0 (15–160)
<i>Mus</i> spp.	13.7 (8–23)
<i>Grammomys</i> spp.	70.2 (55–92)
<i>Dasymys</i> spp.	116.6 (not provided)
<i>Oenomys hypoxantus</i>	110.8 (74–138)
<i>Lamottemys okuensis</i>	71.8 (62–85)
<i>Hybomys</i> spp.	63.7 (55–75)
<i>Lemniscomys</i> spp.	41.1 (28–67)
<i>Otomys occidentalis</i>	77 (69–88)
<i>Atherurus africanus</i>	2,950 (2,500–3,400)
<i>Cricetomys</i> spp.	998.6 (550–1,465)
Squirrels	280.1 (12–770)

Reference

MONADJEM, A., TAYLOR, P.J., DENYS, C., COTTERILL, F.P.D. (2015) *Rodents of Sub-Saharan Africa: a Biogeographic and Taxonomic Synthesis*. De Gruyter, Berlin, Germany.