

- 1 **Supplementary Table S1. *T*- values and *P*-values of the linear regression model (lm) analysis**
- 2 **of the larval positions at death in behavioural assay 2. Values are given for each comparison**
- 3 **of treatments. N = sample size; d. f. = degrees of freedom.**

	Gr 1	Gr 2	Gr 3	Gr 4	C _b	C _a
Gr 1 (N = 59)	---					
Gr 2 (N = 60)	<i>T</i> = 0.822 <i>P</i> = 0.412	---				
Gr 3 (N = 59)	<i>T</i> = 1.012 <i>P</i> = 0.312	<i>T</i> = 0.195 <i>P</i> = 0.846	---			
Gr 4 (N = 60)	<i>T</i> = 0.950 <i>P</i> = 0.343	<i>T</i> = 0.129 <i>P</i> = 0.898	<i>T</i> = 0.067 <i>P</i> = 0.947	---		
C _b (N = 58)	<i>T</i> = 1.745 <i>P</i> = 0.082	<i>T</i> = 0.934 <i>P</i> = 0.351	<i>T</i> = 0.737 <i>P</i> = 0.461	<i>T</i> = 0.807 <i>P</i> = 0.420	---	
C _a (N = 59)	<i>T</i> = 5.906 <i>P</i> < 0.001	<i>T</i> = 6.753 <i>P</i> < 0.001	<i>T</i> = 6.918 <i>P</i> < 0.001	<i>T</i> = 6.881 <i>P</i> < 0.001	<i>T</i> = 7.626 <i>P</i> < 0.001	---
d. f. = 348						

4