

**Web-appendix 4: Additional results.**

**A. Adequately concealed trials (N=10)**

SM						
-	ST					
-	-0.56 -1.41, 0.33	OPI				
-	-0.38 -1.01, 0.27	0.18 -0.42, 0.76	ET			
-	<b>-1.19</b> <b>-2.26, -0.09</b>	-0.64 -1.60, 0.35	-0.81 -1.67, 0.09	EMDR		
-	-0.57 -1.41, 0.28	-0.02 -0.74, 0.70	-0.19 -0.74, 0.37	0.61 -0.20, 1.42	CT	
-	-0.59 -1.38, 0.25	-0.03 -0.65, 0.60	-0.20 -0.69, 0.31	0.60 -0.16, 1.36	-0.02 -0.47, 0.47	CBT

**B. Large-sized trials (N=7)**

SM						
-	ST					
-	0.24 -0.89, 1.38	OPI				
-	-0.14 -0.76, 0.51	-0.38 -1.42, 0.67	ET			
-	-	-	-	EMDR		
-	-	-	-	-	CT	
-	-0.24 -0.89, 0.38	-0.49 -1.46, 0.47	-0.10 -0.63, 0.39	-	-	CBT

**Web-appendix 4:** Effect sizes and credibility intervals of pair-wise comparisons based on the subgroup of adequately concealed trials (A) and large-sized trials (B). Negative effect sizes indicate superiority of the row defining intervention as compared with the column defining intervention. CBT, cognitive behavioural therapy; CT, cognitive therapy; EMDR, eye movement desensitization and reprocessing; ET, exposure therapy; OPI,

other psychological intervention; SM, stress management; ST, supportive therapies; WL, wait-list.