

Psychomotor Dance Therapy Intervention in Dementia

Supplementary Table S4. Meta-Analysis for Mood items across participants

Participants	Baseline (Days)	Item	Phase A Hi	Phase A Low	Phase B Hi	Phase B Low	PAND Re- scaled	Phi	Confidence Intervals
A (ob)	42	Lack of Energy	37	31	31	20	-0.04	-0.064	-0.235 – 0.115
B	42	Lack of energy	32	33.5	33.5	16	-0.16	-0.188	-0.353 – 0.005
C	42	Depressed appearance	32	14.5	14.5	65	0.54	0.506	0.337 – 0.644
D	34	Depressed appearance	23	23.5	23.5	49	0.21	0.170	-0.009 – 0.342
E	35	Low Motor activity	30	19.5	19.5	49	0.34	0.321	0.142 – 0.481
F (ob)	35	None							
G	28	Depressed appearance	9	13	13	77	0.53	0.265	0.068 – 0.466
H	28	None							
I (ob)	28	None							
J	21	Emotional response	20	9	9	67	0.66	0.571	0.377 – 0.719

(ob) observer; WINPEPI aggregated result: Phi 0.24 (95% C.I.: 0.17 – 0.30) See Supplementary Table S3 for interpretation