

PC4 Comfort Database PID 28361

Data Exports, Reports, and Stats

All data (all records and fields)


Hospital name *(hospital_name)*

Total Count (N)	Missing*
30	3 (9.1%)

Does your CVICU routinely screen for delirium using a validated tool? *(does_your_cvicu_routinely)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	3


Counts/frequency: Always - all patients are screened (15, 45.5%), Sometimes - some patients are screened (12, 36.4%), Never - no patients are ever screened (6, 18.2%)



Which scale does your CVICU use to screen for delirium? *(which_scale_does_your_cvic)*

Total Count (N)	Missing*	Unique
32	1 (3.0%)	4

Counts/frequency: CAPD - Cornell Assessment of Pediatric Delirium (22, 68.8%), pCAM-ICU - Pediatric Confusion Assessment Method ICU (3, 9.4%), psCAM-ICU -Preschool Confusion Assessment Method ICU (0, 0.0%), Other (1, 3.1%), None (6, 18.8%)




If Other, please list which one(s) below *(which_scale_does_your_cvic2)*

Total Count (N)	Missing*
1	32 (97.0%)

When screened, how frequently do you score patients? *(when_screened_how_frequent)*

Total Count (N)	Missing*	Unique
28	5 (15.2%)	3

Counts/frequency: **Once daily** (3, 10.7%), **Twice daily** (24, 85.7%), **Q8 hours** (1, 3.6%), **Q4-6 hours** (0, 0.0%), **Q1-3 hours** (0, 0.0%)




Are delirium scores verbally reported to providers (i.e., discussed on rounds)?

(are_delirium_scores_report)

Total Count (N)	Missing*	Unique
31	2 (6.1%)	2


Counts/frequency: Yes (20, 64.5%), No (11, 35.5%)



If Yes, how frequently? *(if_yes_how_frequently)*

Total Count (N)	Missing*	Unique
20	13 (39.4%)	3

Counts/frequency: Once daily (12, 60.0%), Twice daily (7, 35.0%), Q8 hours (0, 0.0%), Q4-6 hours (0, 0.0%), Q1-3 hours (0, 0.0%), After each assessment is performed (1, 5.0%)




How often do postoperative pediatric cardiac surgical patients experience mixed delirium in your CVICU? *(postop_mixed_delirium)*

Total		

Count (N)	Missing*	Unique
32	1 (3.0%)	3


Counts/frequency: >80% of the time (0, 0.0%), 50-80% of the time (2, 6.3%), < 50% of the time (6, 18.8%), Unknown/not measured (24, 75.0%)



How often do postoperative pediatric cardiac surgical patients experience hypoactive delirium in your CVICU? *(postop_hypoactive)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: >80% of the time (0, 0.0%), 50-80% of the time (0, 0.0%), < 50% of the time (6, 18.2%), Unknown/not measured (27, 81.8%)



How often do postoperative pediatric cardiac surgical patients experience hyperactive delirium in your CVICU? *(postop_delirium)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	4


Counts/frequency: >80% of the time (1, 3.0%), 50-80% of the time (2, 6.1%), < 50% of the time (5, 15.2%), Unknown/not measured (25, 75.8%)



Within the past 12 months, has your nursing team received specific education about delirium screening and/or delirium management? *(within_the_past_12_months)*

Total Count (N)	Missing*	Unique
32	1 (3.0%)	2


Counts/frequency: Yes (19, 59.4%), No (13, 40.6%)



Within the past 12 months, has your medical team received specific education about delirium screening and/or delirium management? *(past_12_months_medical)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Yes (16, 48.5%), No (17, 51.5%)



Does your CVICU team incorporate parent/caregiver observations about their child's behavior when diagnosing delirium? *(does_your_cvicu_team_incor)*

Total Count (N)	Missing*	Unique
32	1 (3.0%)	3


Counts/frequency: **Always** (5, 15.6%), **Sometimes** (25, 78.1%), **Never** (2, 6.3%)



Do you have a delirium management protocol, guideline or pathway? *(do_you_have_a_delirium_man)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2

Counts/frequency: **Yes** (9, 27.3%), **No** (24, 72.7%)




Does your unit consult Psychiatry to aid in the diagnosis and/or management of delirium? *(does_your_unit_consult_psy)*

Total Count (N)	Missing*	Unique

Total Count (N)	Missing*	Unique
33	0 (0.0%)	3

Counts/frequency: **Always** (3, 9.1%), **Sometimes** (21, 63.6%), **Never** (9, 27.3%)




Does your unit consult Neurology to aid in the diagnosis and/or management of delirium?

(does_your_unit_consult_psy_2)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: **Always** (0, 0.0%), **Sometimes** (7, 21.2%), **Never** (26, 78.8%)



Who screens your patients for delirium? Please check all that apply. *(who_screens_your_patients)*

Total Count (N)	Missing*	Unique
29	4 (12.1%)	5


Counts/frequency: **CVICU attending physician** (8, 27.6%), **CVICU nurse practitioner** (7, 24.1%), **CVICU bedside nurse** (27, 93.1%), **Neurology attending physician** (1, 3.4%), **Neurology nurse practitioner** (0, 0.0%), **Psychiatry attending physician** (4, 13.8%), **Psychiatry nurse practitioner** (0, 0.0%), **Clinical psychologist** (0, 0.0%)



After screening is performed, who confirms diagnoses your patients with delirium (decides therapies)? Please check all that apply. *(after_screening_is_perform)*

Total Count (N)	Missing*	Unique
29	4 (12.1%)	6

Counts/frequency: CVICU attending physician (25, 86.2%), CVICU nurse practitioner (17, 58.6%), CVICU bedside nurse (9, 31.0%), Neurology attending physician (0, 0.0%), Neurology nurse practitioner (0, 0.0%), Psychiatry attending physician (12, 41.4%), Psychiatry nurse practitioner (1, 3.4%), Clinical psychologist (3, 10.3%)




Download image

What are typical delirium prevention/treatment strategies? Please choose all that apply.

(what_are_typical_delirium)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	10

Counts/frequency: None (0, 0.0%), Re-orientation (29, 87.9%), Early ambulation (26, 78.8%), Preverbal child/infant parental holding/comfort (31, 93.9%), De-intensification (e.g., discontinuing intravenous lines, discontinuing or minimizing opioids, anticholinergics and benzodiazepines) (33, 100.0%), Promoting day/night cycle (33, 100.0%), Exposure to natural light (29, 87.9%), Sleep hygiene (e.g., sleep/rise same time daily; quiet, relaxing, and dark environment) (30, 90.9%), Minimize night-time interventions (28, 84.8%), Prescribe anti-psychotic agents (31, 93.9%), Other (2, 6.1%)




If Other, please list *(if_other_please_list)*

Total Count (N)	Missing*
2	31 (93.9%)

When prescribed, what antipsychotic medication is most commonly used as the first line agent for delirium management? *(when_prescribed_what_antip)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	5


Counts/frequency: Never prescribed (0, 0.0%), Quetiapine (9, 27.3%), Haloperidol (0, 0.0%), Risperidone (15, 45.5%), Chlorpromazine (0, 0.0%), Olanzapine (4, 12.1%), Lorazepam (1, 3.0%), Midazolam (0, 0.0%), Melatonin (4, 12.1%), Zolpidem (0, 0.0%)



What medication is most commonly prescribed for delirium patients with prolongation of the QT interval (hereditary or acquired)? *(what_medication_is_most_co)*

Total Count (N)	Missing*	Unique
32	1 (3.0%)	7

Counts/frequency: Quetiapine (4, 12.5%), Haloperidol (0, 0.0%), Risperidone (6, 18.8%), Chlorpromazine (0, 0.0%), Olanzapine (4, 12.5%), Lorazepam (1, 3.1%), Midazolam (1, 3.1%), Melatonin (10, 31.3%), Zolpidem (0, 0.0%), None (6, 18.8%)




Do you believe there are long-term complications associated with pediatric delirium?

(do_you_believe_there_are_1)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	1

Counts/frequency: Yes (33, 100.0%), No (0, 0.0%)




Does your CVICU unit allow 24-hour parental presence at the bedside?

(does_your_unit_allow_24_ho)

--	--	--

Total Count (N)	Missing*	Unique
33	0 (0.0%)	1


Counts/frequency: **Yes** (33, 100.0%), **No** (0, 0.0%)



Does your CVICU unit use written daily schedules at the bedside for patients with diagnosis of delirium? *(daily_schedules)*

Total Count (N)	Missing*	Unique
32	1 (3.0%)	3


Counts/frequency: **Always** (2, 6.3%), **Sometimes** (18, 56.3%), **Never** (12, 37.5%)



Does your CVICU employ a formal ABCDEF bundle (Awakening and Breathing Coordination, Delirium monitoring and management, Early mobilization, and Family)? *(does_your_unit_employ_a_fo)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	3


Counts/frequency: **Always** (2, 6.1%), **Sometimes** (1, 3.0%), **Never** (30, 90.9%)



Does your CVICU have an early ambulation protocol? *(does_your_cvicu_have_an_ea)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: **Yes** (11, 33.3%), **No** (22, 66.7%)



Does your CVICU utilize pharmacologic sleep aids to promote sleep for patients during their CVICU stay? *(does_your_cvicu_utilize_ph)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	1


Counts/frequency: **Always** (0, 0.0%), **Sometimes** (33, 100.0%), **Never** (0, 0.0%)



What is the most commonly prescribed sleep aid used in your CVICU? *(what_is_the_most_commonly)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	1


Counts/frequency: Lorazepam (0, 0.0%), Midazolam (0, 0.0%), Melatonin (33, 100.0%), Zolpidem (0, 0.0%)



What nonpharmacological interventions does your CVICU routinely use to assist with acquired circadian rhythm disorders? Check all that apply. *(what_nonpharmacological_in)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	10


Counts/frequency: None (2, 6.1%), Eyeshades /masks/covers (9, 27.3%), Ear plugs/muffs (11, 33.3%), Familiar objects (27, 81.8%), Massage (6, 18.2%), Acupressure (2, 6.1%), Acupuncture (3, 9.1%), Music (21, 63.6%), Healing touch (13, 39.4%), Background or soothing sound machines (29, 87.9%)



Does the CVICU team round with Neurology and Psychiatry consultants (minimum: weekly)? *(does_the_cvicu_team_round)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: **Yes** (1, 3.0%), **No** (32, 97.0%)



Does your cardiac program provide any specific follow-up services (such as neurodevelopmental clinic or outpatient psychiatry/psychology) for children who demonstrated delirium during their recovery from cardiac surgery? *(does_your_cardiac_program)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: **Yes** (9, 27.3%), **No** (24, 72.7%)



What is the physical layout of the unit you work on? *(what_is_the_physical_layout)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: **All private rooms** (25, 75.8%), **All semi-private rooms** (0, 0.0%), **Open multi-bed spaces (ie. pods)** (0, 0.0%), **Combination of any of the above** (8, 24.2%)



Does every patient bed space have a window? *(does_every_patient_bed_spa)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: **Yes** (29, 87.9%), **No** (4, 12.1%)



Does your CVICU have a written protocol for light exposure (e.g., dimming lights for quiet time and during nights) and/or day/night cycling? *(light_exposure)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2

Counts/frequency: Yes (3, 9.1%), No (30, 90.9%)




Does the team open the shades/blinds to let natural light in during daytime hours?

(does_the_team_open_the_sha)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2

Counts/frequency: Yes- always (14, 42.4%), Sometimes (19, 57.6%), Never (0, 0.0%)




Does your CVICU have a written noise reduction protocol? *(noise_reduction_protocol)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2

Total Count (N)	Missing*	Unique
32	1 (3.0%)	2


Counts/frequency: Yes (3, 9.4%), No (29, 90.6%)



Does your CVICU have a written pain management protocol, guideline or pathway for all CVICU patients (i.e., medical and surgical intubated patients)? *(pain_management_protocol)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Yes (13, 39.4%), No (20, 60.6%)



If Yes, what are compliance rates for all CVICU patients? *(if_yes_what_are_compliance)*

Total Count (N)	Missing*	Unique
13	20 (60.6%)	3


Counts/frequency: >80% of the time (6, 46.2%), 50-80% of the time (3, 23.1%), < 50% of the time (4, 30.8%)



Does your CVICU have a written pain management protocol, guideline or pathway utilized for all intubated patients (medical and surgical post-operative) or are there different/distinct protocols for patient subpopulations? *(pain_management_protocol_2)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Yes (9, 27.3%), No (24, 72.7%)



If Yes, what are compliance rates for target populations? *(if_yes_what_are_compliance_2)*

Total Count (N)	Missing*	Unique
9	24 (72.7%)	3


Counts/frequency: >80% of the time (4, 44.4%), 50-80% of the time (4, 44.4%), < 50% of the time (1, 11.1%)



Does your CVICU routinely screen for pain needs? *(screen_for_pain_needs)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	1


Counts/frequency: Always - all patients are screened (33, 100.0%), Sometimes - some patients are screened (0, 0.0%), Never - no patients are ever screened (0, 0.0%)



Which scale does your CVICU use to assess pain? *(pain_scale)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	4

Counts/frequency: FLACC Scale (31, 93.9%), Wong-Baker Faces Pain Scale (19, 57.6%), Numerical Rating Pain Scale (21, 63.6%), Other (5, 15.2%)



If other, please list *(if_other_please_list1)*


Total Count (N)	Missing*
6	27 (81.8%)

On average, how frequently are patients assessed for pain? Please check all that apply.

(scoring_frequency)

Total Count (N)	Missing*	Unique
32	1 (3.0%)	4

Counts/frequency: Once daily (0, 0.0%), Twice daily (0, 0.0%), Q8 hours (0, 0.0%), Q4-6 hours (14, 43.8%), Q1-3 hours (20, 62.5%), Procedure or surgery related (7, 21.9%), PRN (17, 53.1%)




Does your CVICU have a sedation protocol, guideline or pathway for all CVICU patients (i.e., medical and surgical intubated patients)?

(sedation_protocol)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Yes (14, 42.4%), No (19, 57.6%)



If Yes, what are compliance rates for all CVICU patients? *(if_yes_what_are_compliance_3)*

Total Count (N)	Missing*	Unique
14	19 (57.6%)	3


Counts/frequency: >80% of the time (6, 42.9%), 50-80% of the time (6, 42.9%), < 50% of the time (2, 14.3%)



Does your CVICU have a sedation protocol, guideline or pathway for targeted CVICU patients (i.e., medical or surgical intubated patients?) *(sedation_protocol_2)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Yes (10, 30.3%), No (23, 69.7%)



If Yes, what are compliance rates for targeted CVICU patients? *(if_yes_what_are_compliance_4)*

Total Count (N)	Missing*	Unique
10	23 (69.7%)	3


Counts/frequency: >80% of the time (2, 20.0%), 50-80% of the time (7, 70.0%), < 50% of the time (1, 10.0%)



Does your CVICU routinely screen for sedation/agitation needs? *(screen_for_sedation_agitation)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	3

Counts/frequency: Always - all patients are screened (30, 90.9%), Sometimes - some patients are screened (2, 6.1%), Never - no patients are ever screened (1, 3.0%)




Which scale does your CVICU use to screen for sedation/agitation needs?

(scale_for_sedation_agitation)

Total Count (N)	Missing*	Unique
31	2 (6.1%)	3


Counts/frequency: State Behavior Scale (23, 74.2%), Comfort (4, 12.9%), RASS (7, 22.6%)



When screened, how frequently do you routinely assess patients for agitation and sedation? (*scoring_frequency_sedation*)

Total Count (N)	Missing*	Unique
32	1 (3.0%)	3


Counts/frequency: Once daily (0, 0.0%), Twice daily (0, 0.0%), Q8 hours (0, 0.0%), Q4-6 hours (11, 34.4%), Q1-3 hours (19, 59.4%), PRN (2, 6.3%)



Are patient sedation goals set and discussed on rounds? (*are_patient_sedation_goals*)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Yes (28, 84.8%), No (5, 15.2%)



How often do postoperative pediatric cardiac surgical patients experience iatrogenic withdrawal syndrome (IWS) at your institution? *(iws_freq)*

Total Count (N)	Missing*	Unique
32	1 (3.0%)	2


Counts/frequency: >80% of the time (0, 0.0%), 50-80% of the time (6, 18.8%), < 50% of the time (26, 81.3%)



Does your CVICU routinely screen pediatric postoperative cardiac surgery patients for IWS? *(iws_screen)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Always (19, 57.6%), Sometimes (14, 42.4%), Never (0, 0.0%)



If always or sometimes, what triggers screening for IWS? *(if_sometimes_what_triggers)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	4


Counts/frequency: Duration of medications (17, 51.5%), Peak dose of medications (0, 0.0%), Cumulative dose of medications (1, 3.0%), Provider discretion (5, 15.2%), At time of weaning protocol (10, 30.3%)



Does your CVICU use a specific withdrawal screening scale? *(does_your_cvicu_use_a_spec)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2

Counts/frequency: Yes (32, 97.0%), No (1, 3.0%)




Which scale does your unit use for withdrawal? *(which_scale_does_your_unit)*

Total Count (N)	Missing*	Unique
32	1 (3.0%)	1

Counts/frequency: WAT-1 (Withdrawal Assessment Tool-1) (32, 100.0%), SOS (Sophia Observation Scale) (0, 0.0%),

Other (0, 0.0%)




When screening for withdrawal, how frequently do you routinely score patients?

(when_screening_for_withdra)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	4

Counts/frequency: Once daily (0, 0.0%), Twice daily (15, 45.5%), Q8 hours (3, 9.1%), Q4-6 hours (14, 42.4%), Q1-3 hours (1, 3.0%), PRN (0, 0.0%)




When a patient is diagnosed with IWS, how frequently do you score them for withdrawal?

(when_a_patient_is_diagnose)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	4

Counts/frequency: Once daily (0, 0.0%), Twice daily (7, 21.2%), Q8 hours (2, 6.1%), Q4-6 hours (19, 57.6%), Q1-3


hours (5, 15.2%), PRN (0, 0.0%)



What is your first line medication for treatment of opioid withdrawal? *(what_is_your_first_line_me)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	3


Counts/frequency: Methadone (20, 60.6%), Morphine (12, 36.4%), Clonidine (1, 3.0%)



What is your first line medication for treatment of opioid withdrawal for patients with evidence of prolonged QT interval (hereditary or acquired)? *(opioid_withdrawal_qt)*

Total Count (N)	Missing*	Unique
32	1 (3.0%)	3


Counts/frequency: Methadone (8, 25.0%), Morphine (20, 62.5%), Clonidine (4, 12.5%)



What is your first line medication for benzodiazepine withdrawal? *(bzd_withdrawal)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Lorazepam (27, 81.8%), Diazepam (6, 18.2%), Clonazepam (0, 0.0%), Midazolam (0, 0.0%), Clonidine (0, 0.0%)



What is your first line medication for alpha agonist withdrawal? *(bzd_withdrawal_2)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Clonidine (31, 93.9%), Dexmedetomidine (2, 6.1%)



Does your CVICU have a process or pathway for distinguishing opioid vs benzodiazepine vs alpha-agonist withdrawal? *(pathway_for_distinguishing_opioid_vs_benzodiazepine_vs_alpha_agonist_withdrawal)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Yes (2, 6.1%), No (31, 93.9%)



Does your CVICU have a written protocol, guideline or pathway that addresses duration of withdrawal medication therapy? *(pathway_that_addresses_duration_of_withdrawal_medication_therapy)*

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2

Counts/frequency: Yes (14, 42.4%), No (19, 57.6%)




If Yes, how often is it used as written? *(if_yes_how_often_is_it_use)*

Total Count (N)	Missing*	Unique

17 | [16 \(48.5%\)](#) | 3


Counts/frequency: >80% of the time (5, 29.4%), 50-80% of the time (7, 41.2%), < 50% of the time (5, 29.4%)



Is there a clinical pharmacist used in the management of opioid withdrawal medication duration? (*is_there_a_clinical_pharma*)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2


Counts/frequency: Yes (30, 90.9%), No (3, 9.1%)



Complete? (*pc4_comfort_database_complete*)

Total Count (N)	Missing*	Unique
33	0 (0.0%)	2

Counts/frequency: Incomplete (1, 3.0%), Unverified (0, 0.0%), Complete (32, 97.0%)



* Note: Values listed as 'Missing' may include records with a Missing Data Code (if Missing Data Codes are defined).