

Table S2. Odds ratios (95% CIs) from logistic regression models for maternal HDP and child motor or mental development abnormalities at the age of 3 years, stratified by gestational weeks.

Variables	No. of cases	No. of cases with HDP		model 1 ^a		model 2 ^b		model 3 ^c	
	n	n	(%)	OR (95% CI)	p-value	OR (95% CI)	p-value	OR (95% CI)	p-value
Motor abnormality									
< 37 weeks	59	6	(10.2)	1.06 (0.45-2.50)	0.893	0.86 (0.35-2.12)	0.738	0.89 (0.37-2.14)	0.801
≥ 37 weeks	365	13	(3.6)	1.69 (0.97-2.95)	0.065	1.54 (0.88-2.70)	0.133	1.60 (0.91-2.81)	0.099
Mental abnormality									
< 37 weeks	28	4	(14.2)	1.57 (0.54-4.56)	0.408	1.44 (0.47-4.43)	0.525	1.35 (0.45-4.05)	0.593
≥ 37 weeks	519	23	(4.4)	2.14 (1.40-3.27)	<0.001	1.92 (1.25-2.94)	0.003	1.96 (1.28-3.00)	0.002

The reference group was defined as children born from mothers without HDP in each group.

^aUnadjusted model.

^bMultivariable logistic models were adjusted for maternal age, multiple pregnancy, parity, maternal smoking, maternal drinking, sex of the child, asphyxia, jaundice, and small for gestational age.

^cMultivariable logistic models were adjusted for maternal age, multiple pregnancy, parity, maternal smoking, maternal drinking, sex of the child.

OR, odds ratio; CI, confidence interval; HDP, hypertensive disorders of pregnancy.