

Supplementary table 2. Sociodemographic and clinical variables at baseline between women who remained in and women who left the study at 8 weeks and 12 months

	Participated at 8 weeks (n = 121)	Did not participate at 8 weeks (n = 10)	Statistical test	Participated at 12 months (n = 112)	Did not participate at 12 months (n = 19)	Statistical test
SES ^a factor score, mean (SD)	0.121	-0.256	$U = 464.0, z = -1.222, p = 0.222$	0.129	-0.121	$U = 875.0, z = -1.235, p = 0.217$
Parity, primiparous, n (%)	63 (52.1)	3 (30.0)	$\chi^2(2) = 2.343, p = 0.310$	60 (53.6)	6 (31.6)	$\chi^2(2) = 4.981, p = 0.083$
Antidepressant usage in pregnancy, yes, n (%)	21 (17.4)	4 (40.0)	$\chi^2(2) = 3.067, p = 0.080$	22 (19.6)	3 (15.8)	$\chi^2(2) = 0.156, p = 0.693$
BDI ^b score at baseline, mean (SD)	9.57 (10.51)	15.72 (17.11)	$t(121) = 1.173, p = 0.266$	9.83 (11.46)	11.92 (10.38)	$t(121) = 0.707, p = 0.481$
STAI-S ^c score at baseline, mean (SD)	36.92 (14.43)	43.91 (16.97)	$t(121) = 1.510, p = 0.134$	37.09 (14.70)	40.37 (15.09)	$t(121) = 0.853, p = 0.396$
Number episodes of MDD prior to pregnancy, >2, n (%)	40 (33.3)	5 (55.6)	$\chi^2(2) = 2.318, p = 0.314$	40 (31.0)	5 (26.3)	$\chi^2(2) = 0.917, p = 0.632$
Maternal history of childhood maltreatment, yes, n (%)	46 (41.8)	5 (50)	$\chi^2(2) = 0.251, p = 0.616$	44 (41.9)	7 (46.7)	$\chi^2(2) = 0.122, p = 0.727$

^aSocioeconomic Score

^bBeck Depression Inventory

^cState-Trait Anxiety Inventory- State